

JP4.17 TORNADO AND SEVERE WEATHER CLIMATOLOGY AND PREDICTABILITY BY ENSO PHASE IN THE NORTH CENTRAL U.S.: A COMPOSITING STUDY

Barbara E. Mayes^{*1}

NOAA/NWS WFO Quad Cities IA/IL, Davenport, Iowa

Craig Cogil

NOAA/NWS WFO Des Moines, Johnston, Iowa

Glenn R. Lussky and Jeffrey S. Boyne

NOAA/NWS WFO La Crosse, Wisconsin

Richard S. Ryhrholm

NOAA/NWS WFO Sioux Falls, South Dakota

1. INTRODUCTION

Climatological factors, including the phase of the El Niño-Southern Oscillation (ENSO), are known predictors for variables such as seasonal temperature and precipitation and tropical cyclone activity. ENSO phase also has a relationship to severe weather activity and may have utility in predicting anomalies in severe weather climatology, including the number of significant tornadoes and the number of tornado days. While not useful in daily severe weather operations and forecasting, determining the relationship between ENSO phase and severe weather climatology anomalies can aid preparedness for potentially active seasons among operational meteorologists, emergency managers, the media, and others with a role in preparing spotters and citizens for severe weather seasons.

Previous studies have investigated the relationship between ENSO and tornado climatology in the United States and Canada, with varying definitions of ENSO and methodologies (Bove and O'Brien, 1998; Browning 1998; Etkin et al., 2001; Wikle and Anderson, 2003). Agee and Zurn-Birkhimer (1998) determined that an axis of increased tornado activity during La Niña years extended from Iowa through Illinois and Indiana into Kentucky and Tennessee, while an axis of increased tornado activity during El Niño years extended from Colorado and New Mexico through the Texas panhandle into Oklahoma and Missouri. They concluded that their findings were a result of geographical shifts in tornado activity, rather than an overall increase or decrease in activity nationwide based on ENSO phase. Bove (1998) found a similar axis of increased activity in La Niña years.

The north central United States is the focus of this study, encompassing nine states (North Dakota, South Dakota, Nebraska, Kansas, Minnesota, Wisconsin, Iowa, Missouri, and Illinois) in the central and northern

Plains and middle and upper Mississippi River valley that are climatologically in the heart of activity during the spring and summer convective seasons. Data are grouped by National Weather Service (NWS) Weather Forecast Office (WFO) County Warning Area (CWA) to capture smaller-scale climatological features.

This severe weather compositing study will serve many purposes. Primarily, the study is an investigation of the impact of a large-scale climatological factor that exhibits predictability, ENSO, on a smaller-scale weather phenomenon, severe convection. Embedded within this study, as well, is a case study of the methodology of compositing that will be used operationally by the NWS as the 3-Month Outlook of Local El Niño/La Niña Impacts (3MOLEI). For this purpose, examining data at the CWA level will demonstrate the utility of the method for NWS WFO weather and climate operations.

2. DATA AND METHODOLOGY

Conducting a statistical compositing study of severe weather climatology based on ENSO requires a consistent and approved definition of ENSO as well as a quality data set and a rigorous compositing methodology.

2.1 ENSO Definition

The National Oceanic and Atmospheric Administration (NOAA) developed an operational definition for ENSO utilizing the Oceanic Niño Index (ONI). ONI is defined as the three-month running average sea surface temperature anomaly in the Niño 3.4 region. An El Niño occurs when ONI is at least 0.5 °C for at least five consecutive 3-monthly periods, or “seasons”; conversely, La Niña conditions occur when ONI reaches -0.5 °C or less for five consecutive seasons.

¹ Corresponding author address: Barbara E. Mayes, National Weather Service, 9050 Harrison St., Davenport, IA, 52806; e-mail: Barbara.Mayes@noaa.gov

The ONI has been calculated by 3-month period and is available online via NOAA's Climate Prediction Center (CPC) back to 1950 (Smith and Reynolds, 2003). The NOAA definition of ENSO based on ONI is followed in this study.

2.2 Data Source

Tornado and severe weather data for this study were collected from NOAA's National Climatic Data Center (NCDC) Storm Data publication, in both its online database and document formats. Other quality data sets do exist (e.g., Grazulis 1993), but their interpretation in relation to the NCDC records can be subjective and can introduce a source of inconsistency. In addition, Grazulis (1993) provides data for significant tornadoes only and not the full tornado record.

The NOAA Storm Data publication does have known limitations, including the upward trend in reports due to increasing population coverage, more aggressive verification practices, and an increase in the number of storm chasers (Verbout et al., 2006). Data missing from the online Storm Data database in June and July 1993 were added manually based on the Storm Data documents. NCDC Storm Data were filtered by the authors to remove reports that did not meet severe criteria (hail 0.75 inches or greater; wind 50 knots or greater) as well as wind reports that were non-convective. Tornadoes in the NCDC database are listed by county; thus, to prevent multiple counts of the same tornado, county border-crossing tornadoes were grouped together as one tornado. Finally, several instances of duplicate reports, particularly during the early 1990s, were removed. The resulting database of well over 150,000 storm reports provides a clean foundation for the study, albeit with the inherent limitations described above.

While the number of tornado reports does show an upward trend with time, particularly in the number of F0 and F1 tornadoes, the number of tornado days and the number of significant (F2 or greater) tornadoes does not show nearly as much of a trend. Therefore, the number of significant tornadoes was used for the entire period of record of ENSO phase (1950-2005), and the number of tornado days was used from 1960-2005. The hail database exhibits a strong upward trend in the number of reports well into the 1990s, even when looking at the number of significant (2 inch or greater) hail days (Doswell et al., 2005); thus, the conditional climatologies using hail records to ENSO phase likely are contaminated by that trend. The wind database was even more inconsistent due to changes in reporting style through the years as winds were estimated, listed at 0 knots, or listed with wind speeds not available when direct measurements were not made. As a result, convective winds have not been addressed in this study.

2.3 Compositing Methodology

The compositing technique used in this study was developed in partnership among NWS headquarters, CPC, and NWS staff in regional and field offices. The resulting product will become an operational forecast product, the 3MOLEI, at the NWS WFOs, with the flexibility for offices to composite any weather or climate variable that has a statistically meaningful relationship to the ENSO phase as determined by the methodology. Conditional climatologies will be combined with forecast probability of an ENSO phase to predict the probability of a given climatological variable occurring in the above, near, or below normal category given the ENSO phase that is forecast to occur. While the forecast information will be a critical part of the 3MOLEI, the forecast portion of the compositing methodology is not addressed in this study. Much of the detail about the compositing methodology is available online as training material and templates at:

<http://www.nws.noaa.gov/om/csd/pds/pcu4/IC4.3/index.htm>

Compositing analysis employs a sampling technique based on the conditional probability of a given event occurring based on the ENSO phase. The result is a conditional climatology of a given variable based on the ENSO phase. The methodology incorporates a standardized method for developing climatologies of the given parameter for each ENSO phase (La Niña, neutral, and El Niño) in comparison to the climatology of the period of record, as well as incorporating a test for statistical significance of the results to determine if deviations from the average climatology are significant relative to the null hypothesis that there is no relationship between ENSO phase and each climatological variable.

In order to test for statistical significance, the compositing analysis methodology determines whether there is a significant ($p \leq 0.10$) relationship between ENSO phase and the climatological variable. The probability of a given climatological variable occurring in the above, near, or below normal categories exactly x number of times, $P(x)$, within a given ENSO phase is a hypergeometric distribution. If the number of occurrences, x , falls within the lower 10% or upper 10% of the distribution, the event is deemed statistically significant, and the null hypothesis that the climatological variable occurred in the above, near, or below normal category during a given ENSO phase by chance is rejected.

This study focuses on developing the conditional climatology of several severe weather climatological variables, focusing on the number of tornado days per year and the number of significant (F2+) tornadoes per year. The study considers weak, moderate, and strong episodes of El Niño and La Niña to be part of the climatology in those phases, therefore not differentiating ENSO phase by its strength.

3. RESULTS

Initial results indicate many coherent signals in all phases of ENSO across the study area. Relationships of severe weather climatologies to the ENSO phase were made from the fall prior to the “convective year,” or year from which severe weather data were analyzed (beginning August-September-October of the prior year), through the fall of the convective year (September-October-November). The yearly total of significant tornadoes and tornado days was composited to the ENSO phase for each three-month season, with the purpose of investigating lagged relationships between ENSO phase and convective activity as well as concurrent impacts.

Results typically were found in a given CWA during any one of three periods: fall and winter of the prior year (August-September-October of the prior year through December-January-February), early spring of the convective year (January-February-March through April-May-June), and late spring/summer/fall of the convective year (May-June-July through September-October-November). Due to the spring frailty of the ENSO phase, which is the tendency for the ENSO phase to be more sensitive to change in April and May, the ENSO phase during the early spring seasons

tended to carry a less reliable statistical relationship to convective activity and often represented a shift in the type of impact within a given state or CWA. The results found in this study were consistent with several of the previous studies noted above, including Agee and Zurn-Birkhimer (1998), Bove (1998), Bove and O’Brien (1998), Browning (1998), and Etkin (2001).

In Figures 1 and 2, results were shaded if they appeared in more than one season within the period (prior year fall/winter or convective year late spring/summer/fall), ensuring temporal consistency and further reducing the potential for chance to influence results. On the rare occasions when the significant tornado (ST) and tornado day (TD) signals were opposed—for example, if the signal during a La Niña favored a diminished number of significant tornadoes but an enhanced number of tornado days—the results were not plotted.

In both periods detailed below, individual CWAs demonstrated statistically significant relationships across multiple seasons without coherent signals in neighboring areas. The compositing analysis was created to determine relationships at the WFO scale, and due to smaller scale influences as well as the potential for smoothed or conflicting signals within




Figure 1. Impacts on significant tornado (ST) and tornado day (TD) climatologies during the late spring, summer, and early fall of the concurrent convective year based on ENSO phase. Unshaded sections within the study area exhibited either no statistically significant signal or conflicting signals.

neighboring CWAs, results within individual CWAs are not required to match neighboring CWAs for the results to be useful. For example, the Green Bay, WI, (GRB) WFO exhibited an enhanced potential for a below normal number of significant tornadoes when neutral conditions occur during the late spring/summer/fall period (Figure 1). Occurrences such as that in GRB demonstrate the benefit of the compositing methodology within a CWA in determining local scale statistical relationships to the ENSO phase.

3.1 Late Spring/Summer/Fall

A wide swath of the study area demonstrated a tendency for enhanced tornado activity when a La Niña is occurring in the late spring/summer/fall of the concurrent convective year, with relationships to both the number of tornado days and the number of significant tornadoes (Figure 1). Statistically significant signals appeared consistently for either an enhanced probability of above normal activity or a diminished probability of below normal activity during a La Niña across much of Illinois and Missouri into Kansas and Nebraska, as well as along the northern tier areas of Wisconsin, Minnesota, and North Dakota. The relationship was among the strongest observed in the study area for all ENSO phases and seasons for consistency among seasons as well as spatial

consistency. Some of the CWAs within the region also exhibited diminished potential for tornado activity during either El Niño or neutral phases, including a signal for enhanced potential of below normal tornado activity during neutral years in Iowa and Illinois as well as Nebraska into eastern Kansas and western South Dakota.

Conversely, in central Wisconsin and Minnesota, signals for diminished tornado activity during a La Niña appeared, as well as signals for enhanced potential for activity during either El Niño or neutral phases extending into South Dakota.

3.2 Prior Year Fall/Winter

While results were not as coherent as those for ENSO phases occurring during the heart of convective season, several significant signals did appear based on the ENSO phase of the fall and winter prior to the convective year (Figure 2). In an area centered on Nebraska, signals indicate an enhanced probability of below normal tornado activity when an El Niño occurs during the preceding fall and winter. The signal also appears, though less spatially consistent, in parts of Wisconsin and Missouri.

Relationships also appeared for the La Niña phase. In




Figure 2. Impacts on significant tornado (ST) and tornado day (TD) climatologies during the fall and winter prior to the convective year based on ENSO phase. Unshaded sections within the study area exhibited either no statistically significant signal or conflicting signals.

southeastern Minnesota and Wisconsin, signals appeared for enhanced potential of below normal tornado activity or diminished potential of above normal tornado activity when a La Niña occurs during the preceding fall and winter. Parts of Nebraska and Kansas, as well as the southeastern study area in Missouri and Illinois, exhibited signals for enhanced potential of above normal or diminished potential of below normal activity when a La Niña occurs in the preceding fall and winter, as well.

4. CONCLUSIONS AND FUTURE WORK

The results of this study indicate that there are statistically significant relationships between multiple severe weather parameters and ENSO phase in the north central United States. Results were attainable in areas with a high number of severe weather events as well as those where severe weather occurrences are less frequent, indicating that the compositing methodology will be useful for severe weather parameters even in areas outside of the peak of severe weather activity.

Extensive future work will be required before broad conclusions can be drawn. The study will be expanded to include more severe weather parameters, and composites may be completed to compare activity within a three-month season to the ENSO phase, rather than comparing yearly activity to the phase, to better determine temporal as well as spatial patterns in the relationship between ENSO and severe weather activity. Statistical results will need to be translated to synoptic pattern changes for given seasons and ENSO phases, as well, in order determine relationships between ENSO phase and anomalies in severe weather climatology in the north central United States.

Acknowledgments

This study was greatly aided by the work of several colleagues and students, including Ray Wolf (WFO Quad Cities IA/IL), Benjamin R. J. Schwedler (Iowa State University/WFO La Crosse, WI), Jeff Erickson (Texas A&M/WFO Quad Cities IA/IL), and Jennifer Laflin (University of Nebraska-Lincoln/WFO Sioux Falls, SD). The compositing methodology was developed by Marina Timofeyeva (NWS HQ/UCAR), Robert Livezey (NWS HQ), and Annette Hollingshead (NWS), with contributions in both the methodology and training by several others, including Heather Hauser (NWS) and Andrea Bair (NWS). Adnan Akyuz (NWS) made valuable contributions to the manuscript.

REFERENCES

Agee, E., and S. Zurn-Birkhimer, 1998: Variations in USA tornado occurrences during El Niño and La Niña. *Preprints, 19th Conf. on Severe Local Storms, AMS*, 287-290.

Bove, M. C., 1998: Impacts of ENSO on United States tornadic activity. *Preprints, 19th Conf. on Severe Local Storms, AMS*, 313-316.

Bove, M. C., and J. J. O'Brien, 1998: Impacts of ENSO on United States tornadic activity. *Preprints, 9th Symposium on Global Change Studies, AMS*, 199-202.

Browning, P., 1998: ENSO related severe thunderstorm climatology of northwest Missouri. *Preprints, 19th Conf. on Severe Local Storms, AMS*, 291-292.

Doswell, C. A., H. E. Brooks, and M. P. Kay, 2005: Climatological estimates of daily local nontornadic severe thunderstorm probability for the United States. *Wea. Forecasting*, **20**, 4, 577-595.

Etkin, D., S. E. Brun, A. Shabbar, and P. Joe, 2001: Tornado climatology of Canada revisited: tornado activity during different phases of ENSO. *Intl. J. Clim. (Royal Meteorological Society)*, **21**, 915-938.

Grazulis, T. P., 1993: *Significant Tornadoes, 1680-1991*. Environmental Films, 1326 pp.

Smith, T. M., and R. W. Reynolds, 2003: Extended reconstruction of global sea surface temperatures based on COADS data (1854-1997). *J. Climate*, **16**, 10, 1495-1510.

Verbout, S. M., H. E. Brooks, L. M. Leslie, and D. M. Schultz, 2006: Evolution of the U.S. tornado database: 1954-2003. *Wea. Forecasting*, **21**, 1, 86-93.

Wikle, C. K., and C. J. Anderson (2003): Climatological analysis of tornado report counts using a hierarchical Bayesian spatio-temporal model. University of Missouri, available online at http://www.stat.missouri.edu/~wikle/TORNADOpap_v3.pdf