


NATURE AND CHARACTERISTICS OF WEAK AND INTENSE KATABATIC FLOWS

Jon A. Arrillaga ⁽¹⁾, Carlos Yagüe ⁽¹⁾, Carlos Román-Cascón ^(1,2), Mariano Sastre ⁽¹⁾, Jordi Vilà-Guerau de Arellano ⁽³⁾, Gregorio Maqueda ⁽¹⁾

(1) DEPT. FÍSICA DE LA TIERRA Y ASTROFÍSICA, UNIVERSIDAD COMPLUTENSE DE MADRID, SPAIN (jonanarr@ucm.es) (2) LABORATOIRE D'AÉROLOGIE, UNIVERSITY OF TOULOUSE, CNRS, FRANCE (3) METEOROLOGY AND AIR QUALITY GROUP, WAGENINGEN UNIVERSITY, NETHERLANDS

1. LA HERRERÍA

This analysis is performed in *La Herrería Forest*, located at the foothill of the Guadarrama Mountain Range (Spain), at around 50 km from the city of Madrid.



Summers are particularly dry and warm in this region. The surface energy balance is therefore moisture limited!

2. OBSERVATIONS



In this work we use 10-minutal meteorological measurements carried out during an **intensive summer campaign** in 2017 (22/06 – 26/09).

Picture of the 10-m meteorological fixed tower at La Herrería Forest during Summer 2017. La Herrería is part of the Guadarrama Monitoring Network (GUMNet; www.ucm.es/gumnet).

Main measurements used in this study:

Variable	Height (m, agl)	Instrument
Air temperature	3, 6, 10	Aspirated thermometer
Wind speed	3, 6, 10	Cup anemometer
Wind direction	10	Wind Vane
Turbulent fluxes	4, 8	IRGASON
Rain	surface	Pluviometer
Soil Moisture	-0,05	Reflectometer
Soil-heat flux	-0,05	Heat-flux plate
Radiation components	2	4-component radiometer

3. KATABATIC DETECTION



We apply an objective and systematic algorithm to the observational data in order to select the events that fulfil **katabatic-event criteria**:

Filter	Criteria	Description
1	Weak large-scale winds	$V_{850} < 6 \text{ m s}^{-1}$
2	Days without synoptic cold fronts	$(\Delta\theta_{e,850}/\Delta t) > -1.5^\circ\text{C}/6\text{h}$
3	Non-rainy events	$pp < 0.5 \text{ mm}/\text{day}$
4	Minimum persistence in the katabatic direction	$WD \in [240 - 330]_{2h}$

It is based on the algorithm for selecting sea-breeze events from Arrillaga et al. (2018).

We find 40/94 katabatic events meeting the requirements from the algorithm.


Wind rose at 6 m over 2 h after the katabatic onset.



1. WE CLASSIFY KATABATICS INTO 3 GROUPS ACCORDING TO THEIR MAXIMUM INTENSITY AT 6 M.

WHICH ARE THE FACTORS PRODUCING DIFFERENT INTENSITIES?

MOTIVATION




2. KATABATIC INTENSITY IS ASSOCIATED WITH DISTINCT REGIMES IN THE STABLE BOUNDARY LAYER (SBL)

HOW DOES THE INTERACTION BETWEEN KATABATICS AND TURBULENCE OCCUR?

4. NATURE

4.1) Close relationship between katabatic intensity & thermal stratification at the onset




THERMAL STRATIFICATION: $\Delta\theta_v = \theta_v(10\text{m}) - \theta_v(3\text{m})$
 Unstable: $(\Delta\theta_v)_{\text{onset}} < -0.2 \text{ K}$
 Neutral: $-0.2 \text{ K} \leq (\Delta\theta_v)_{\text{onset}} \leq +0.2 \text{ K}$
 Stable: $(\Delta\theta_v)_{\text{onset}} > +0.2 \text{ K}$

Early onset (prior to 2h before sunset) → Unstable stratification → Intense (& few rest) katabatics
 Late onset (after 2h before sunset) → Neutral/Stable stratification → Weak (& most rest) katabatics

4.2) Thermal stratification at the katabatic onset is directly linked to turbulence.

Maximum katabatic intensity (between onset and 00 UTC) at 6 m vs turbulent kinetic energy (TKE) at 8 m at the onset




WE FIND TWO GROUPS:
 (1) $TKE \sim 0.1 \text{ m}^2 \text{ s}^{-2}$ → Neutral/Stable stratification (WEAK AND SOME REST KATABATICS)
 (2) $TKE \sim 1 \text{ m}^2 \text{ s}^{-2}$ → Unstable stratification (INTENSE AND SOME REST KATABATICS)

TURBULENCE INTENSITY AT THE ONSET MODULATES THE SUBSEQUENT KATABATIC INTENSITY → IT IS EXPLAINED BY THE COMPLEX INTERACTION BETWEEN KATABATICS AND TURBULENCE

4.3) Which are the factors that induce an earlier or later onset of katabatic flows?

Maximum katabatic intensity (colours) at 6 m & NCEP-reanalysis wind direction at 850 hPa.



We explore the influence of synoptic-wind direction and soil moisture (Fitzjarrald, 1984; Banta & Gannon, 1995).

Reanalysis wind speed at 850 hPa for some events is indicated with white labels to show its possible influence.


Since it is generally weak, its influence is negligible

N-NW SYNOPTIC WIND + SOIL MOISTURE $< 0.06 \text{ m}^3 \text{ m}^{-3}$ → INTENSE KATABATICS
 W-SW / S-SE SYNOPTIC WIND or N-NW SYNOPTIC WIND + SOIL MOISTURE $> 0.06 \text{ m}^3 \text{ m}^{-3}$ → WEAK KATABATICS

5. INTERACTION WITH TURBULENCE

5.1) Downward sensible heat flux and surface energy balance

Downward sensible-heat flux ($H < 0$) vs the katabatic intensity at 6 m (left) and the heat-flux demand (HFD = $-(Q_N - G)$) at surface (right).



- $H < 0$ is maximum for intermediate wind speed (1.5 – 3.5 m s^{-1}).
- We neglect the latent-heat flux (it is almost zero) in the surface energy balance and compare H vs HFD.
- The greatest negative energetic imbalance occurs for weak katabatics, which induces the formation of a very stable regime.

5.2) Regime transition from non-dimensional parameters


We characterise the regime transition from relevant non-dimensional parameters by representing thermal stratification $\Delta\theta_v$ and V_{TKE} as a function of one local and one non-local parameter: z/L and shear capacity (SC) respectively.

$$V_{TKE} = \sqrt{\frac{1}{2}(u'^2 + v'^2 + w'^2)}$$

This variable is used as a measure of turbulence intensity.

$$L = \frac{-u_*^3 \theta_v}{kgw \theta_v'}$$

Turbulent fluxes are calculated at 8 m agl and so is z/L .



By using this local parameter we are unable to find a single value to separate the SBL regimes!

For instance, results are similar to Mahrt et al. (1998) at 10 m for the regime transition:
 → Weakly stable: 0.1 VS 0.065 (Mahrt et al., 1998)
 → Very stable: 0.5 VS 1 (Mahrt et al., 1998)


The SC compares the measured shear with the minimum shear to maintain continuous turbulence, which is given by the heat-flux demand at surface (van Hooijdonk et al., 2015). We calculate it considering that turbulence depends on local shear when $U < V_g \approx 1.5 \text{ m s}^{-1}$, and on bulk shear when $U > V_g$ (HOST transition).

$$SC(U < 1.5 \text{ m s}^{-1}) = \frac{\rho C_p (\kappa z)^2 (\partial U / \partial z)^{3/3}}{g / \theta_v |Q_N - G|}$$

$$SC(U > 1.5 \text{ m s}^{-1}) = \frac{\rho C_p (\kappa z)^2 (U / z)^{3/3}}{g / \theta_v |Q_N - G|}$$

Local shear is calculated from fitting wind-speed measurements to a log-linear profile. We consider a minimum HFD of 50 W m^{-2} to sustain continuous turbulence.

Except for very few 10-min values, weak katabatics are linked with the very stable SBL and intense katabatics with the weakly stable SBL.



This non-local parameter predicts the regime transition ($SC \sim 3.5$) independently of the height (van Hooijdonk et al., 2015).

THIS PARAMETER IS DEFINED BY IGNORING TRANSPORT TERMS IN THE TKE BUDGET, WHICH HOWEVER CAN BE IMPORTANT IN KATABATIC WINDS. STILL, WE ARE ABLE TO DEFINE THE REGIME TRANSITION FROM THE KATABATIC WIND SPEED.

6. FINAL THOUGHT

WE HAVE LINKED THE SYNOPTIC WIND AND LOCAL SOIL-MOISTURE CONDITIONS, WITH KATABATIC INTENSITY AND THE ASSOCIATED SBL REGIMES FROM THEIR INTERACTION WITH TURBULENCE:

SYNOPTIC WIND IN THE KATABATIC DIRECTION + LOW SOIL MOISTURE → EARLY KATABATIC ONSET: UNSTABLE/CONVECTIVE STRATIFICATION → MODERATE TO HIGH TURBULENCE → BULK SHEAR INCREASES → SUBSEQUENT INTENSE KATABATIC → NEAR-NEUTRAL/WEAKLY STABLE SBL REGIME.

IF THE CONDITIONS ABOVE FOR THE SYNOPTIC WIND AND SOIL MOISTURE ARE NOT MET → LATER ONSET: NEUTRAL/STABLE STRATIFICATION → WEAK TURBULENCE + RADIATIVE LOSS NOT COMPENSATED → STABLE STRATIFICATION THAT SUPPRESSES TURBULENCE PRODUCTION (POSITIVE FEEDBACK) → WEAK KATABATIC PRODUCING A VERY STABLE REGIME.

7. REFERENCES

Arrillaga, J., Vilà-Guerau de Arellano, J., Bosveld, F., Baltink, H., Yagüe, C., Sastre, M., and Román-Cascón, C., 2018: Impacts of afternoon and evening sea-breeze fronts on local turbulence, and CO_2 and radon-222 transport. *Q. J. R. Meteorol. Soc.*, in press. DOI: 10.1002/qj.3252.

Banta, R. M. and Gannon, P. S. T., 1995: Influence of soil moisture on simulations of katabatic flow. *Theor. Appl. Climatol.* 52, 85–94.

Fitzjarrald, D. R., 1984: Katabatic Wind in Opposing Flow. *J. Atmos. Sci.*, 41, 1143–1158.

Mahrt, L., 1998: Nocturnal Boundary-Layer Regimes. *Boundary-Layer Meteorol.*, 88, 255–278.

Sun, J., Mahrt, L., Banta, R. M., and Pichugina, Y. L., 2012: Turbulence Regimes and Turbulence Intermittency in the Stable Boundary Layer during CASES-99. *J. Atmos. Sci.*, 69, 338–351.

van Hooijdonk, I. G. S., Donda, J. M. M., Clercx, H. J. H., Bosveld, F. C., and van de Wiel, B. J. H., 2015: Shear Capacity as Prognostic for Nocturnal Boundary Layer Regimes. *J. Atmos. Sci.*, 72, 1518–1532.

8. ACKNOWLEDGEMENTS

This research has been funded by the ATMOUNT-II project [Ref. CGL2015-65627-C3-3R (MINECO/FEDER)] and the Project [Ref. CGL2016-81828-REDT/AEI] from the Spanish Government, and by the GUMNet (Guadarrama Monitoring Network, www.ucm.es/gumnet) observational network of the CEI Moncloa campus of International Excellence. Jon A. Arrillaga is supported by the Predoctoral Training Program for No-Doctor Researchers of the Department of Education, Language Policy and Culture of the Basque Government (PRE_2017_2_0069, MOD = B). We thank the contribution of all the members of the GUMNet Team, and Patrimonio Nacional for the facilities given during the installation of the meteorological tower.

