



Radar-based rainfall nowcasting at European scale


long-term evaluation and performance assessment

Marc Berenguer, Daniel Sempere-Torres











OPERA radar mosaic:

Precipitation
 observations over
 Europe @2 km and
 every 15 minutes.




OPERA radar mosaic:

- Precipitation
 observations over
 Europe @2 km and
 every 15 minutes.
- Operationally produced since mid 2011.



OPERA radar mosaic:

- Precipitation
 observations over
 Europe @2 km and
 every 15 minutes.
- Operationally produced since mid 2011.



OPERA radar mosaic The European NEXRAD?

The European NEXRAD?

Many C-band radars.

- Many C-band radars.
- OPERA network is extremely heterogeneous:

- Many C-band radars.
- OPERA network is extremely heterogeneous:
 - -installation date,

- Many C-band radars.
- OPERA network is extremely heterogeneous:
 - -installation date,
 - -manufacturers,

- Many C-band radars.
- OPERA network is extremely heterogeneous:
 - -installation date,
 - -manufacturers,
 - -scanning strategy,

- Many C-band radars.
- OPERA network is extremely heterogeneous:
 - -installation date,
 - -manufacturers,
 - -scanning strategy,
 - -signal processing and QC,

- Many C-band radars.
- OPERA network is extremely heterogeneous:
 - -installation date,
 - -manufacturers,
 - -scanning strategy,
 - -signal processing and QC,
 - -and product generation.


- Many C-band radars.
- OPERA network is extremely heterogeneous:
 - -installation date,
 - -manufacturers,
 - -scanning strategy,
 - -signal processing and QC,
 - -and product generation.
- OPERA radar density is about twice that of NEXRAD.


- Many C-band radars.
- OPERA network is extremely heterogeneous:
 - -installation date,
 - -manufacturers,
 - -scanning strategy,
 - -signal processing and QC,
 - -and product generation.
- •OPERA radar density is about twice that of NEXRAD.





Frequency of rain (dBZ > 10 dBZ)


in the period 01 June - 30 June 2013







































The HAREN project

The goal:


Demonstrate the interest of *rainfall*nowcasting based on OPERA mosaics for hazard assessment.







Very short-term forecasting by extrapolation of radar observations.



Very short-term forecasting by extrapolation of radar observations.

I.Tracking: The motion field of rainfall is estimated from observations.


Very short-term forecasting by extrapolation of radar observations.

I.Tracking: The motion field of rainfall is estimated from observations.




Very short-term forecasting by extrapolation of radar observations.

I.Tracking: The motion field of rainfall is estimated from observations.




Very short-term forecasting by extrapolation of radar observations.

- I.Tracking: The motion field of rainfall is estimated from observations.
- 2.Extrapolation: Of the most recent rainfall field.



Very short-term forecasting by extrapolation of radar observations.

- I.Tracking: The motion field of rainfall is estimated from observations.
- **2.Extrapolation:** Of the most recent rainfall field.








How does it work?

-performance of nowcasting-









Dbservations 13 July 2012 21:45 UTC


Forecasts




Observed

Forecasted









Evolution of the life time in the period 01 June - 31 July 2012

Lifetime mostly between 4 and 10 h




Evolution of the life time in the period 01 June - 31 July 2012

Lifetime mostly between 4 and 10 h




Evolution of the life time in the period 01 June - 31 July 2012


Lifetime mostly between 4 and 10 h




Mean Lifetime in the period 01 June - 31 July 2012



Mean Lifetime in the period 01 June - 31 July 2012




Mean Lifetime in the period 01 June - 31 July 2012




Case #1: 13 June 2012 1200 UTC

nowcasts up to 8 hours




Case #2: 25 June 2012 0715 UTC


nowcasts up to 8 hours




Geographical variability (Lifetime)

in the period 01 June - 31 July 2012







Long-term evaluation in the period 01 June - 31 May 2013





Long-term evaluation in the period 01 June - 31 May 2013




Long-term evaluation in the period 01 June - 31 May 2013














When is nowcasting useful?


-comparison with NWP outputs-









01 June 2013 - 48 hour evaluation




01 June 2013 - 48 hour evaluation





01 June 2013 - 48 hour evaluation







01 June 2013 - 48 hour evaluation







01 June 2013 - 48 hour evaluation







01 June 2013 - 48 hour evaluation







01 June 2013 - 48 hour evaluation







01 June 2013 - 48 hour evaluation







01 June 2013 - 48 hour evaluation






01 June 2013 - 48 hour evaluation







01 June 2013 - 48 hour evaluation







01 June 2013 - 48 hour evaluation






01 June 2013 - 48 hour evaluation







•Continental nowcasts benefit from the better capturing of motion at large scale and upstream coverage.

- Continental nowcasts benefit from the better capturing of motion at large scale and upstream coverage.
- Nowcasting shows average lifetime ~5 hours. Very similar to the results obtained over the NEXRAD mosaic.

- Continental nowcasts benefit from the better capturing of motion at large scale and upstream coverage.
- Nowcasting shows average lifetime ~5 hours. Very similar to the results obtained over the NEXRAD mosaic.
- Time and space variability:

- Continental nowcasts benefit from the better capturing of motion at large scale and upstream coverage.
- Nowcasting shows average lifetime ~5 hours. Very similar to the results obtained over the NEXRAD mosaic.
- Time and space variability:
 - Large scale systems are more predictable.

- Continental nowcasts benefit from the better capturing of motion at large scale and upstream coverage.
- Nowcasting shows average lifetime ~5 hours. Very similar to the results obtained over the NEXRAD mosaic.
- Time and space variability:
 - Large scale systems are more predictable.
 - ▶ Better skill in the areas with more frequent precipitation.

- Continental nowcasts benefit from the better capturing of motion at large scale and upstream coverage.
- Nowcasting shows average lifetime ~5 hours. Very similar to the results obtained over the NEXRAD mosaic.
- Time and space variability:
 - Large scale systems are more predictable.
 - ▶ Better skill in the areas with more frequent precipitation.
- •First comparisons against operational NWP show cross-over times ~4-5 hours.







Radar-based rainfall nowcasting at European scale long-term evaluation and performance assessment



Spring 2008 over North-America





From: Berenguer et al. (MWR2012)

OU Storm-Scale Ensemble Forecast (10 members).

- WRF-based with perturbed configurations.
- 4-km grid.
- 9 members with 3D-var assimilation of reflectivity and Doppler velocity.