



Early Kinematic and Thermodynamic Structural Differences Between Developing and Non-developing Atlantic Tropical Convective Systems

Charles N. Helms (chip.helms@gmail.com) and Robert E. Hart (rhart@fsu.edu)

Department of Earth, Ocean, and Atmospheric Sciences, Florida State University, Tallahassee, Florida



Introduction

- Numerous studies have detailed how the structure of a developing tropical convective system evolves (e.g. Bister and Emanuel 1997)
- Comparatively fewer studies have examined structural differences between developing and non-developing systems (e.g. McBride and Zehr 1981)
 - Majority of these use composite means
- The present study aims to observationally examine individual cases of developing and non-developing systems in hopes of identifying key differences in the vorticity structure early in the evolution of the system.

Important Signatures of Tropical Cyclogenesis

- Vorticity magnitude is greater, on average, in developing systems (McBride and Zehr 1981)
- Formation of an **upright vortex** (e.g. Bister and Emanuel 1997, Simpson et al. 1997, Ritchie and Holland 1997, Montgomery et al. 2006)
- High **moisture** and strong mid-level circulation (Nolan 2007)
- Vortex alignment, moist mid-levels, strengthening mid-level circulation (Davis and Ahijevych 2012)

Data and Methodology

- ### Data
- GRIP and PREDICT dropsondes
 - Time-space correction using mean zonal wind at each level

Method (as per Helms and Hart 2012)

- Vorticity calculated using Green's Theorem
- Polygonal regions used for calculations (see Fig. 1)

$$\text{Vorticity} \approx \frac{\sum(u\Delta x + v\Delta y)}{\text{area}}$$




Fig. 1. Visual outline of method for calculating vorticity using Green's Theorem on polygonal regions (from Helms and Hart 2012).

Developing Case - PGI44L - Karl (2010)




Fig. 2. Series of cross-sections (earlier times on left) from the developing Pre-Karl system which was declared a tropical depression on Sept. 14 at 12 UTC. For each time, plots of the corresponding IR satellite image (top row, left), 500 hPa relative vorticity (top row, middle), 800 hPa relative vorticity (top row, right), relative vorticity cross-section (middle row), and relative humidity cross-section (bottom row) are included. The red or black horizontal bars in the satellite image and the 800 hPa and 500 hPa relative vorticity plots represent the location of the cross-section data at the corresponding time. The dashed line in the vorticity and relative humidity cross-sections indicates the approximate axis of maximum vorticity.

Developing Case - PGI46L - Matthew (2010)




Fig. 3. As in Fig. 2, except for the developing Pre-Matthew system which was declared a tropical depression on Sept. 23 at 12 UTC.

Acknowledgments

This work is funded through the NASA Genesis and Rapid Intensification Processes (GRIP) Grant #NNX09AC43G. Data support for the GRIP and PREDICT field experiments is provided by NCAR/EOL under sponsorship of the National Science Foundation.

Non-Developing Case - PGI27L




Fig. 4. As in Fig. 2, except for the non-developing PGI27L system.

Conclusions

With respect to the three systems analyzed here:

- Vorticity structure evolves from a highly tilted to a **vertically aligned vorticity column in both developing and non-developing systems**
- Although not shown here, no significant differences were found in divergence and temperature anomaly structures in relation to the vorticity structure
- Only noticeable structural difference is in the presence of **dry air at mid-levels in the non-developing case**
 - Confirms findings of Nolan (2007) and Davis and Ahijevych (2012)
- These results suggest that the **key determinant of genesis in these cases was the co-location of moisture with the column of maximum vorticity**

References

Bister, M., and K. A. Emanuel, 1997: The genesis of Hurricane Guillermo: TEXMEX analyses and a modeling study. *Mon. Wea. Rev.*, **125**, 2662–2682.

Davis, C. A., and D. A. Ahijevych, 2012: Mesoscale structural evolution of three tropical weather systems observed during PREDICT. *J. Atmos. Sci.*, **69**, 1284–1305.

Helms, C. N., and R. E. Hart, 2012: A polygon-based line integral method for calculating vorticity, divergence, and deformation from non-uniform observations. *Extended Abstracts, 30th Conf. on Hurricanes and Tropical Meteorology*, Ponte Vedra, Florida, Amer. Meteor. Soc.

McBride, J. L., and R. Zehr, 1981: Observational analysis of tropical cyclone formation. Part II: Comparison of non-developing versus developing systems. *J. Atmos. Sci.*, **38**, 1132–1151.

Montgomery, M. T., M. E. Nicholls, T. A. Cram, and A. B. Saunders, 2006: A vortical hot tower route to tropical cyclogenesis. *J. Atmos. Sci.*, **63**, 355–386.

Ritchie, E. A., and G. J. Holland, 1997: Scale interactions during the formation of Typhoon Irving. *Mon. Wea. Rev.*, **125**, 1377–1396.

Simpson, J., E. A. Ritchie, G. J. Holland, J. Halverson, and S. Stewart, 1997: Mesoscale interactions in tropical cyclone genesis. *Mon. Wea. Rev.*, **125**, 2643–2661.