

— Is the lady dead, was she killed and by whom? —



Artwork: Michael Schrenk

© von Storch, HZG

An attempt to deconstruct the observed climate trends in the Baltic Sea and Med Sea Basins

Hans von Storch (HZG, OUC),
Armineh Barkhordarian and
Roberto C Mechoso (UCLA)



14 January 2016


28th Conference on Climate Variability and Change, New Orleans, AMS, 12C.3

 Helmholtz-Zentrum
Geesthacht
Zentrum für Material- und Küstenforschung

 Ocean University of China



Fig. 4.11 Annual and seasonal mean surface air temperature anomalies (relative to 1960–1991) for the Baltic Sea basin 1871–2011, calculated from 5° by 5° latitude, longitude box average taken from the CRUTEM3v dataset (Brohan et al. 2006) based on land stations (from top to bottom: a annual, b winter (DJF), c spring (MAM), d summer (JJA), e autumn (SON)). Blue comprises the Baltic Sea basin north of 60°N and red south of 60°N. The dots represent individual years and the smoothed curves (Gaussian filter, $\sigma = 3$) highlight variability on timescales longer than 10 years



Baltic Sea region air temperature development



MANIFESTATION





Observed temperature trends in the Baltic Sea region (1982-2011)

Baltic Sea region




Estimating natural variability:

2,000-year high-resolution regional climate palaeo-simulation (Gómez-Navarro et al, 2013) is used to estimate natural (internal + external) variability.

- An external cause is needed for explaining the recently observed annual and seasonal warming over the Baltic Sea area, except for winter (with < 2.5% risk of error)

2m Temperature in the Med Sea Region (1980-2009)

Observed changes of 2m temperature (1980-2009) in comparison with GS signals



■ Observed trends of 2m temperature (1980-2009)

■ 90% uncertainty range of observed trends, derived from 10,000-year control simulations



➤ There is less than 5% probability that natural (internal) variability is responsible for the observed annual and seasonal warming in the Med Sea region, except in winter.

DETECTION





Observed and projected temperature trends in the Baltic Sea Region (1982-2011)




- **DJF and MAM changes can be explained by dominantly GHG driven scenarios**
- **None of the 10 RCM climate projections capture the observed annual and seasonal warming in summer (JJA) and autumn (SON).**

Observed and projected temperature trends in the Med Sea Region (1982-2011)




Observed changes of 2m temperature (1980-2009) in comparison with GS signals



➤ In the Med Sea region, the warming can be explained by the A1B scenario of increased GHGs.

Precipitation trends in the Baltic Sea Region (1979-2008)




In **winter (DJF)** none of the 59 segments derived from 2,000 year paleo-simulations yield a positive trend of precipitation as strong as that observed. There is less than 5% probability that observed positive trends in winter be due to natural (internal + external) variability alone (with less than 5% risk).

In **spring (MAM)**, **summer (JJA)** and **Annual** trends externally forced changes are not detectable. However observed trends lie within the range of changes described by 10 climate change scenarios, indicating that also in the scenarios a systematic trend reflecting external forcing is not detectable (< 5% risk).

In **autumn (SON)** the observed negative trends of precipitation contradicts the upward trends suggested by 10 climate change scenarios, irrespective of the observed dataset used.



Precipitation in the Med Sea Region (Over land, 1966-2005, CMIP3)





ATtribution


Solar surface radiation in the Baltic Sea Region, 1984-2005



1880-2004 development of sulphur dioxide emissions in Europe (Unit: Tg SO₂). (after Vestreng et al., 2007 in BACC-2 report, Sec 6.3 by HC Hansson)

- A possible candidate to explain the observed deviations of the trends in summer and autumn, which are not captured by 10 RCMs, is the effect of changing regional aerosol loads


Surface solar radiation in the Med Sea Region (1985-2004)



Decrease of anthropogenic aerosols due to:

- ✓ more effective clean-air regulations, energy consumption
- ✓ decline in the Eastern European economy in the late 1980s, closure of dirty factories

Changes in Large-scale circulation (SON) in terms of sea level pressure



- Observed trend pattern shows areas of decrease in SLP over the Med. Sea and areas of increase in SLP over the northern Europe. Observed trend pattern of SLP in SON contradicts regional climate projections.
- **The mismatch between projected and observed precipitation in autumn is already present in the atmospheric circulation.**



ATtribution

Conclusions

- ✓ An influence of non-natural signal is detectable in spring, summer and fall winter for temperature and winter and fall in precipitation changes.
- ✓ The observed temperature changes in all seasons, and in winter and spring in precipitation are in the direction of what scenarios suggest.
- ✓ However there are inconsistencies between observed changes and scenarios.
 - temperature changes are stronger than what scenarios suggest
 - observed precipitation changes in late summer and autumn contradict projected changes.
- ✓ The analysis of large-scale circulation patterns, in terms of mean sea-level pressure and geopotential height at 500 hPa, confirms the inconsistency detected for precipitation.

Conclusion

Our analysis indicates that the recent regional climate change in Europe cannot be explained, in the framework of our present knowledge, without reference to elevated greenhouse gases. However, in summer and fall, the driver „GHGs“ is insufficient in explaining the recent change.

Possible causes:

- a) Suggestion for response to GHG driver by climate model is inaccurate.
- b) Other drivers are significant, in particular the non-maintenance of the earlier atmospheric aerosol-load (Problem: we have no regional quantified guess patterns)
- c) Natural variability is underestimated by historical simulation with climate model - the change is still within the range of natural variability.

Some science is settled, but lots of science is not settled.