

P6.4 POLARIMETRIC PROTOTYPE OF THE WSR-88D RADAR OBSERVATIONS OF BIRDS AND INSECTS

Pengfei Zhang, Alexander Ryzhkov, and Dusan Zrnic*

CIMMS, University of Oklahoma, Norman, Oklahoma

*NOAA/National Severe Storms Laboratory, Norman, Oklahoma

1. INTRODUCTION

Biological echoes are treated as contamination like ground or sea clutters by radar meteorologists as they estimate rainfall intensities or assimilate radar reflectivity into numerical forecast model. Doppler velocity measured by radar in the presence of small insects well represents air motion because small insects are considered as passive tracers. But in the presence of birds especially migrating birds, Doppler velocities measured by radar are different from air velocities (Jungbluth et al. 1995; Gauthreaux et al. 1998a, b).

On the other hand, biological echoes are very useful for ornithologists to study the migration, dispersion pattern and flight behaviors of insects and birds especially during nighttime. As early as 1969, Schaefer (1969) had successfully used radar as an entomological tool to study the flight trajectories of locusts. Now more and more information has been published in the fields of radar entomology (<http://www.ph.adfa.edu.au/a-drake/trews/>) and radar ornithology (<http://virtual.clemson.edu/groups/birdrad/>).

The application of polarimetric technique on weather radar provides more information about biological scatterers. It has been shown that polarimetric radar has capability to recognize birds and insects (Zrnic and Ryzhkov 1998). Azimuthal dependencies of differential reflectivity Z_{dr} observations have been used to determine the occurrence of migrations of birds and insects in the clear air when the radar observed wind was analyzed (Wilson et al 1994). Achtemeier (1991) used local minimums of Z_{dr} to conclude that insects in the updraft region are in evasive flight and cannot be treated as passive tracers. Recently, Lang et al. (2004) reports that CSU-CHILL polarimetric radar observed quasi-symmetric echo patterns of reflectivity and Z_{dr} in clear air.

It is obvious that polarimetric radar become a very useful tool for researchers to learn about the characteristics of different scatterers in the clear air. Dual-polarimetric capability will be implanted on the currently operating WSR-88D weather radar network in the near future. It will benefit not only meteorological studies and weather forecast but also ecology studies.

Clear air observations with the KOUN polarimetric radar in 2004 provide unique opportunity for us to study the spatial distribution and temporal variability of the echo from birds and insects in the planetary boundary layer during daytime and nighttime. The analysis of observations on 8 September 2004 is presented in this paper.

2. OBSERVATIONS ON 8 SEPTEMBER 2004

It is well known that in order to escape from unfavorable living conditions in both space and time and colonize themselves in a region with favorable conditions, insects and birds migrate polewardly in spring and summer and equatorwardly in autumn every year. Different types of synoptic weather provide favorable airflows for the migrations of insects and birds (Drake and Gatehouse 1995; Dingle 1996). Most passerines migrate during night (Corral 1989). These nocturnal migrants fly dispersedly rather than concentrate or narrow their route around land formation like diurnal migrants. Thus, radar often observe relative smooth "disk-like" echo in fall and spring (Gauthreaux et al. 1998a; Zhang et al. 2005).

On 8 September 2004, a high pressure system set up over Oklahoma and slowly moved to the east. Atmosphere was quite stable and north-northeast wind dominated the lower atmosphere. It provided favorable airflows for southwardly migrating birds and insects in the fall season. The KOUN radar continuously operated on 8 September 2004




Fig. 1 KOUN radar observations at 0.5° elevation angle at 06 UTC (upper panels) and 20 UTC (lower panels). Panels (a) and (e) represent reflectivity Z ; (b) and (f) differential reflectivity Z_{dr} ; (c) and (g) differential phase ϕ_{dp} ; (d) and (h) Doppler velocity V_r . Radar is located right at the center of each panel. Two white range rings indicate 50 km and 100 km ranges from the radar.




Fig. 2 Surface weather maps at 12 UTC on (a) 7 September and on (b) 8 September 2005.

and recorded valuable data. Fig.1 shows reflectivity Z (a and e), differential reflectivity Z_{dr} (b and f), differential phase ϕ_{dp} (c and g), and Doppler velocity (d and h) observed with the KOUN radar at 0.5° elevation angle at 06 UTC (upper panels) and 20 UTC (lower panels). It is obvious that the echo coverage is larger in the upper panels (06 UTC) than in the lower panels (20 UTC) except (d) and (h). The average reflectivity Z is also stronger at 06 UTC than 20 UTC. The echoes within 30 km from the radar are attributed to ground clutter. Larger coverage and stronger echoes indicate more scatterers and they reach higher altitude (about 1800 m ASL) at 06 UTC. The obvious visual differences in Z_{dr} and ϕ_{dp} imply the sizes of dominated scatterers are clearly different. The wind direction from Doppler velocity V_r (d

and h) at 06 UTC and 20 UTC indicates northeast wind through lower boundary layer (below 1.5 km) that is consistent with rawinsonde observations and weather pattern shown in the weather map (Fig. 2).

3. EVOLUTIONS OF POLARIMETRIC VARIABLES

Four sectors centered at four different azimuths 30° , 120° , 210° , and 300° at 0.5° elevation angle are selected in order to study the characteristics of radar moments along and perpendicular to the ambient wind direction (Fig. 3). The width of each sector is 30° . The ranges from 40 km to 70 km are selected. The reasons are 1) to avoid contaminations of ground clutter near the




Fig. 3 Schematic diagram for the locations of four sectors (cyan trapezoid). Radar is right at the center of range rings. The gray arrow indicates the direction of ambient wind.

radar; and 2) to gather most available valid observations because clear air echo beyond 70 km in the daytime (see Fig. 1 lower panel) is quite weak.

Average Z , Z_{dr} , and ϕ_{dp} for the four sectors are calculated and the evolution of these variables is illustrated in Fig. 4a, b, and c. Note that straight lines between 15 UTC and 17 UTC in Fig. 4 correspond to the lack of observations. In Fig. 4 all the radar variables show abrupt changes just after sunset and before sunrise. The sunset time is 00:46 UTC and sunrise time is 12:08 UTC on 8 September 2004. For example, average Z at azimuth 210° quickly drops to -13 dBZ after sunset and then climbs back and reaches its maximum of 22 dBZ in about 1 hour. Just before sunrise, it suddenly drops to its minimum -18.4 dBZ at 11:46 UTC and bounced back to 15.4 dBZ at 12:03 UTC. The abrupt changes of radar variables at sunset and sunrise in the clear air day during migrating season are common. Similar pattern is also found in the observations from the KOUN radar on 8-9 May and 12 June 2004 (not shown). It implies that the birds and insects dramatically change their flight behavior as the sky light changes.

Radar ornithology studies (Lincoln et al. 1998) have found that nocturnal migrants fly at

different altitudes at different times during the night. Shortly after sunset, migrants generally take off and rapidly reach their maximum altitude. They keep flying at the peak altitude until around midnight and gradually descend until sunrise. Our observations (Fig. 4a) show similar evolution pattern. For the average Z of all four sectors, they rapidly reach their peak in about 1.5 hour after sunset and then gradually decrease.

More information about biological scatterers is provided by polarimetric variables (Fig. 4b and c). Vaughn (1985) suggested that prolate spherical water drop is a better model for birds and insects as radar scatterers than spherical water drops (Riley, 1985). Usually the ratios of width-to-length are between 1:2 and 1:3 for birds and between 1:3 and 1:10 for insects (Vaughn, 1985). Due to the difference in shape and size, the polarimetric signatures of insects and birds are different. Based on the studies of Zrnic and Ryzhkov (1998), as polarimetric radar scatterers birds have lower Z_{dr} and higher ϕ_{dp} than insects. Fig. 1 and 4 indicate that Z_{dr} is roughly lower in the nighttime (between 00:46 UTC to 12:08 UTC) and ϕ_{dp} is lower in the daytime (between 12:08 UTC to 24:00 UTC). Hence, dominant scatterers are birds in the nighttime and insects in the daytime. Note that we use term “dominant scatterers” here. Actually we found




Fig. 4 Time evolution of (a) reflectivity Z , (b) differential reflectivity Z_{dr} , and (c) differential phase ϕ_{dp} for the four sectors at four different azimuths 30° (green), 120° (red), 210° (blue), and 300° (pink).

that birds and insects frequently coexist in a radar resolution volume (Bachmann and Zrnica 2005).

Fig. 4 also demonstrates that the spatial and temporal variations of average Z_{dr} and ϕ_{dp} are obviously larger than average Z especially in the nighttime. Average Z for all four sectors (Fig.4a) gradually decreases with time from their peak value at about 02:00 UTC in the nighttime. However, average Z_{dr} and ϕ_{dp} show distinct diurnal differences among four sectors. This is likely because Z_{dr} and ϕ_{dp} are independent of the concentration of scatterers but are dramatically affected by their size, shape and orientation (Doviak and Zrnica 1993). Also reflectivity is less sensitive to the size change of scatterers than Z_{dr} and ϕ_{dp} (Zrnica and Ryzhkov 1998).

The strong azimuthal variations of Z_{dr} and ϕ_{dp} are primarily attributed to the orientation of biological scatterers. Variation in the scatterers size and type is another possible cause of diversity. The latter argument is consistent with conclusions made in some of biological studies. Lewis and Taylor (1965) have studied about 400 species of insects and found that different species fly at different times of the day or night. Ornithologists found that average migration distances for different birds vary from 50 km/day to 110 km/day when they migrate across northern America (Dingle 1996). Assuming that average flight speed relative to the ground is 15 m/s, it only takes about 1 to 2 hours of continuous flight for individual bird to finish its one day migratory journey.

4. AZIMUTHAL DEPENDENCE

Due to preferred orientation of biological scatterers, their signatures would exhibit a well-pronounced azimuthal dependence if they migrate in the same direction with respect to radar. The azimuthal dependencies of Z_{dr} from insects were observed by Mueller and Larkin (1985), Achtemeier (1991) and Lang et al. (2004). Zrnic and Ryzhkov (1998) have compared the azimuthal dependencies of Z_{dr} and ϕ_{dp} for reflections from birds with model results for oblate spheroids. The model results showed good qualitative agreement with the observations. The important result of that study is that Z_{dr} and ϕ_{dp} reach their maxima as the angle θ between the direction of incident radar beam and the symmetry axis of prolate spheroid is equal to 90° . In the Rayleigh regime of scattering typical for small insects, Z_{dr} and ϕ_{dp} monotonically increase to their minima at $\theta = 90^\circ$. In the Mie regime of scattering typical for large insects and birds, azimuthal dependences of Z_{dr} and ϕ_{dp} are more complicated. But their values at $\theta = 0^\circ$ are still smaller than at $\theta = 90^\circ$.

Polarimetric observations (Fig. 4) do not exhibit azimuthal dependence predicted by the model. During the nighttime, ϕ_{dp} is obviously larger at azimuth 120° (red line) than at other three azimuths. According to the model, ϕ_{dp} at azimuth 300° (pink line) should be similar to the one at azimuth 120° . But ϕ_{dp} at azimuth 300° is much smaller than at azimuth 120° . Z_{dr} exhibits more complicated pattern during the nighttime. Before 06 UTC, the pattern of Z_{dr} at four different directions is similar except Z_{dr} at azimuth 30° (green line) is smaller than the others. Between 06 UTC to 10 UTC, Z_{dr} at azimuth 300° becomes the largest among the four directions. There are two reasons to explain why pronounced azimuthal dependence is not observed. First, different birds with different sizes and shapes migrate at the same time. Second, the mixture ratio of birds and insects may be different at different sectors. We speculate that high Z_{dr} and relative low ϕ_{dp} at azimuth 300° may indicate more insects are in the sector between 06 UTC and 10 UTC.

Between 18:00 UTC and 23:00 UTC, Z_{dr} in the sectors perpendicular to the ambient wind direction (azimuths 120° and 300°) are clearly smaller than in the sectors along the wind

direction (azimuths 30° and 210°). Oval-shaped radar echo pattern is observed (Fig. 1e) during that period of time. As we mentioned previously, the dominant scatterers in the daytime are insects. Large Z_{dr} and strong echo along the ambient wind direction imply the orientation of insects is perpendicular to the wind direction. Common orientation of migrating insects, such as grasshoppers and moths, has been observed and studied by entomologists. The angle between the orientation direction of insects and ambient wind direction is defined as "crab angle" by Wolf et al. (1995). Generally, crab angle is less than 90° . But in this case, the crab angle is about 90° . It is different from the observation of Lang et al. (2004). They found that the insects are flying upwind, so the crab angle is 180° in that case. It is also different from migrating birds. They usually take the advantage of tail-wind and their orientation direction is the same as ambient wind direction.

5. COMPARISON BETWEEN RADAR MEASURED WIND AND SOUNDING

The ambient wind directions at 1000 m ASL observed by rawinsonde are 40° and 48° at 00 UTC and 12 UTC separately. Thus, the V_r observations in the sectors centered at 30° and 210° are compatible with the rawinsonde observed wind speed. KOUN radar is collocated with the OUN sounding observation station. With north-northeast wind, sounding balloon drifted to south-southwest of the radar as it ascends. The azimuth of 210° is closer to the balloon than the azimuth of 30° . Time evolution of average V_r , Z , and Z_{dr} in the sector at azimuth 210° is illustrated in Fig. 5. The wind speeds from rawinsonde observations at 00 UTC and 12 UTC at height 1000 m ASL that is about same as the height of the KOUN radar beam at range 55 km are also marked as blue circles in Fig. 5. The ambient wind directions are slightly different from radar azimuth 210° . Thus the wind speeds shown in Fig. 5 are the projections of rawinsonde observations to the azimuth 210° . They are 11.3 m/s and 9.2 m/s at 00UTC and 12 UTC. The sunset (00:46 UTC) and sunrise (12:08 UTC) time are also marked with brown dash lines in Fig. 5.

Average V_r slowly increases from 11.3 m/s (the same as rawinsonde observation at 00:00




Fig. 5 Time evolution of average Z (red), Z_{dr} (green), and V_r (blue) of the sector at azimuth 210° for 8 September 2004. Blue circles indicate rawinsonde observations at 00 UTC and 12 UTC. Brown dash lines indicate sunset and sunrise time.

UTC) to its peak of 17.5 m/s at 06:45 UTC. Then it slowly drops to 10.1 m/s at 11:40 UTC. Average V_r between sunset and sunrise is clearly higher than ambient wind speed observed by rawinsonde at 00 UTC and 12 UTC. Since no other observations rather than Doppler radar are available, true wind speed in the atmosphere is unknown between 00 UTC and 12 UTC. Weather maps (Fig. 2) show the weather pattern does not change much over Oklahoma. We believe that wind speed in the lower atmosphere should not vary much during the night. Hence, the high “wind speed” observed by radar is the sum of ambient wind speed and speed of migrating birds. This conclusion is consistent with the polarimetric observations.

6. NOCTURNAL MIGRATING INSECTS AND BIRDS

The observations of the KOUN radar between 02:00 to 04:00 UTC provide an

excellent example to show the capability of polarimetric radar to distinguish insects and birds as they coexist in the clear air environment. Fig. 6a shows a disk-like reflectivity field observed with the KOUN at 0200 UTC. It is similar to the observations of current WSR-88D radar when migrating birds are present (Gauthreaux et al. 1998a, b). The reflectivity field is relative uniform and smooth. The decline of intensity with range indicates the decrease of concentration of scatterers with height. The smoothness implies that the scatterers are similar. But the observations of polarimetric variables Z_{dr} and ϕ_{dp} clearly reveal that the clear air echoes are not uniform as the reflectivity shows. A small area with local minimum of Z_{dr} and a maximum of ϕ_{dp} are found at range of 50 km and azimuth of 240° (Fig.6b and c). Meanwhile, a large area of high Z_{dr} (> 7 dB) and low ϕ_{dp} ($< 80^\circ$) is around it. High Z_{dr} and low ϕ_{dp} indicate that the dominate scatterers are insects in this area. In contrast, birds are dominated scatterers in the




Fig. 6 KOUN radar observations (a) reflectivity Z , (b) differential reflectivity Z_{dr} , and (c) differential phase ϕ_{dp} at 0.5° elevation angle at 0220 UTC. Solid blue lines indicate the city limit. The red arrow in (a) indicates the location of city Chickasha.

small area. This observation reveals a large flock of insects was migrating in the night. As the city limit (blue lines in Fig. 6) is overlapped onto the radar observations, it is found that the small area with local minimum of Z_{dr} and a maximum of ϕ_{dp} just locates above the city Chickasha. It may imply that birds are favorite to city light and those nocturnal migrating insects avoid the light, or their predators, or both. At the same time, a large flock of birds characterized by low Z_{dr} and high ϕ_{dp} was migrating on the east side of the radar.

The magnitude of average ambient V_r is about 10 m/s. V_r measurements at this azimuth are only a component of the movement of scatterers. Assuming the biological scatterers were flying at the same direction as the ambient wind. The total speed of scatterer movement at 220° , the direction of ambient wind, is obtained as 10.6 m/s. Also assuming the ambient wind at 02:00 UTC does not change much from the 00:00 UTC when rawinsonde observation is available, the ambient wind speed around 600 m above the ground is about 10.7 m/s. It indicates that insects are driven by the ambient wind, but birds were flying around over the city in the ambient wind.




Fig. 7 Doppler velocity V_r observation along the beam at azimuth of 240° and (a) elevation angle of 0.5° and (b) 1.5° .

The Doppler velocity V_r observation confirms the findings inferred from polarimetric variables. Fig. 7 shows V_r along a beam at elevation angle of 0.5° and 1.5° and azimuth of 240° . Between 50 and 55 km where the city locates at, strong perturbations from ambient V_r are measured at the elevation angle of 0.5° .

The area with high Z_{dr} and low ϕ_{dp} was not observed at the elevation angle of 1.5° (not shown). The V_r measurements at the elevation angle of 1.5° (Fig. 7b) show more dispersive than that at 0.5° . The average V_r between 50 and 60 km (corresponding to 1.5 km above the ground) is about 17 m/s. The estimated total velocity is 18.1 m/s faster than ambient wind speed of 15.8 m/s observed by rawinsonde at that height. This result confirms that migrating birds were dominant scatterers at higher level. It also denotes that the insects were flying below a certain height. Berry and Taylor (1968) has found that if temperature is too low (below about 17°C) insects like aphids, may not fly. The rawinsonde observation at 00:00 UTC shows the air temperature decreases from 20.6°C at height of 550 m to 11.6°C at 1480 m. So the atmosphere at the height of 1.5 km is too cold for insects. It may be the reason why insects are not observed by the radar at the elevation angle of 1.5° .

At 03:19 UTC, about 1 hour later, the polarimetric feature of nocturnal migrating insects around Chickasha almost totally disappeared. The average V_r increases from 10 m/s to 12 m/s although the V_r just above Chickasha is still more dispersive than adjacent area (not shown). These observations indicate insects ceased their migration and migrating birds become dominant scatterers in this area.

The observations of nocturnal migrating insects are also supported by biological studies. Drake (1984) and Drake and Farrow (1988) have found that larger insects commonly migrate during night. Several reasons may explain why large insects migrate in nighttime. First, they may avoid their predators like birds. Second, more stable atmosphere and less thermal stress due to lack of solar radiation may save power and provide an optimal conditions for insect migrants (Rainey 1974).

7. DISCUSSION AND SUMMARY

Many radar meteorologists consider insects as passive tracers, so radar-measured winds represent ambient air motion. For example, Russell and Wilson (1997) have suggested that insects can be viewed as "aerial plankton". But other radar meteorologists argue that it may not be true for large insects such as moths and locusts since they can fly at speed of 3 to 5 m/s (Riley et al. 1983; Riley 1974). As migrating insects orient and travel in almost the same direction, systematic bias in the radar-measured wind can be induced (Achte-meier 1991; Lang et al. 2004). The studies conducted by Srygley et al. (1996), and Walker and Riordan (1981) indicate that butterfly can adjust their airspeed and flight directions against drift induced by adverse wind in order to migrate to the destination. Time-compensated sun can be used as a compass in insect migrants to navigate them at a preferred geographical flight direction (Oliveira et al. 1998). Thus, it may not be appropriate to treat all insects as passive tracers when insects, especially large insects, migrate. Checking the azimuthal dependence pattern observed with polarimetric radar becomes an effective way to detect if insects migrate in the same direction.

The observations of diurnal variation of radar variables clearly reflect some behaviors of migrating birds and insects that are in agreement with biological studies. The sensitivity and azimuthal dependence of polarimetric variables provide ample information to distinguish between birds and insects and estimate the orientation of biological scatterers. Z_{dr} and ϕ_{dp} measurements might also give a chance to roughly estimate the size of birds and insects once the distinction between them is made. The observations of the KOUN radar between 02:00 to 04:00 UTC demonstrate the advantages of polarimetric radar over current operational WSR-88D radar to reveal the detailed spatial structure of different flocks of nocturnal migrating insects and birds.

This work shows the potential of a polarimetric prototype of the WSR-88D radar for detection and classification of biological scatterers. Therefore, polarimetric diversity will benefit data quality control and facilitate assimilation of polarimetric radar data onto weather forecasting models. On the other hand, the biological echoes considered as contamination of weather echoes for meteorologists contain useful information for biologists studying flying behavior of insects and birds. Integrated with other observations such as mark-release-recapture experiments and weather forecast, network polarimetric radar observations can provide vital information regarding insects' and birds' density, their destinations and arrival dates even forecast their behavior for ecology management system.

ACKNOWLEDGEMENT

This conference paper was prepared by Pengfei Zhang with funding provided by NOAA/Office of Oceanic and Atmospheric Research under NOAA-University of Oklahoma Cooperative Agreement #NA17RJ1227, U.S. Department of Commerce. The statements, findings, conclusions, and recommendations are those of the authors and do not necessarily reflect the views of NOAA or the U.S. Department of Commerce.

This research is also partially funded by the Federal Aviation Administration (FAA). The views expressed are those of the authors and

do not necessarily represent the official policy or position of the FAA.

The authors would like to thank the dedicated support from the NSSL and CIMMS/University of Oklahoma staff who maintain the KOUN WSR-88D polarimetric radar.

REFERENCES

- Achtemeier, G. L. 1991: The use of insects as tracers for "clear-air" boundary-layer studies by Doppler radar. *J. Atmos. Ocean. Tech.*, **8**, 746-765.
- Bachmann S., and D.S. Zrnic, 2005: Spectral polarimetry identifies and separates mixed biological scatterers. Preprints, 32nd Conference on Radar Meteorology, Albuquerque, NM, American Meteorological Society, CD-ROM, this issue.
- Corral, M., 1989: The world of birds: A Layman's guide to ornithology. *The Globe Pequot Press*, 224pp.
- Dingle, H., 1996: Migration: The biology of life on the move. *Oxford University Press*, 474pp.
- Doviak, R.J., and D. S. Zrnic, 1993: Doppler Radar and Weather Observations. 2nd edition, *Academic Press*, Page 562.
- Drake, V. A., and A. G. Gatehouse, 1995: Insect migration. *Cambridge University Press*, 478pp.
- Drake, V.A., and R.A. Farrow, 1988: The influence of atmospheric structure and motions on insect migration. *Annual Review of Entomology*, **33**, 183-210.
- Gauthreaux, S. A., and C. G. Belser, 1998a: Displays of bird movements on the WSR-88D: patterns and quantification. *Wea. Forecasting.*, **13**, 453-464.
- Gauthreaux, S. A., D. S. Mizarhi and C. G. Belser, 1998b: Bird migration and bias of WSR-88D wind estimates. *Wea. Forecasting*, **13**, 465-481.
- Jungbluth, K., J. Belles, and M. Schumacher, 1995: Velocity contamination of WSR-88D and wind profiler data due to migrating birds. Preprints, 27th Conference on radar meteorology, 9 - 13 August 1995, Vail, Colorado, Amer. Meteor. Soc., 666 - 668.
- Lang, T. J., S. A. Rutledge, and J.L. Stith, 2004: Observations of Quasi-Symmetric Echo Patterns in Clear Air with the CSU-CHILL Polarimetric Radar. *J. Atmos. Ocean. Tech.*, **21**, 1182-1189.
- Lincoln, C. F., S. R. Peterson, and J. L. Zimmerman, 1998: Migration of Birds. <http://www.npwr.usgs.gov/resource/other/data/migratio/migratio.htm#table>
- Mueller, E. A., and R. P. Larkin, 1985: Insects observed using dual-polarization radar. *J. Atmos. Oceanic Technol.*, **2**, 49-54
- Oliveira, E.G., R.B. Srygley, and R. Dudley, 1998: Do Neotropical migrant butterflies navigate by a solar compass? *J. Exp. Biol.* **201**, 3317-3331.
- Rainey, R.C., 1974: Biometeorology and insects flight: some aspects of energy exchange. *Ann. Rev. Entomol.* **19**, 407-439.
- Riley, J.R., 1974: Radar observations of individual desert locusts (*Schistocerca gregaria*) (Forsk.). *Bull. Ent. Res.*, **64**, 19-32.
- Riley, J.R., D.R. Reynolds, and M.J. Farmery, 1983: Observations of the flight behaviour of the armyworm moth, *Spodoptera exempta*, at an emergence site using radar and infra-red optical techniques. *Ecol. Entomol.*, **8**, 395-418.
- Riley, J. R., 1985: Radar cross sections of insects. *Proc. IEEE*, **73**, 228-232.
- Russell, R.W. and J.W. Wilson, 1997: Radar-observed "fine lines" in the optically clear boundary layer: Reflectivity contributions from aerial plankton and its predators. *Boundary Layer Meteorology*, **82**, 235-262.
- Srygley, R.B., E.G. Oliveira, and R. Dudley, 1996: Wind drift compensation, flyways, and conservation of diurnal, migrant

Neotropical Lepidoptera. *Proc. R. Soc. Lond. B* **263**, 1351-1357.

Vaughn, C. R., 1985: Birds and insects as radar targets: A review. *Proc. IEEE*, **73**, 205-227.

Walker, T.J., and A.J. Riordan, 1981: Butterfly migration: Are synoptic-scale wind system important? *Ecol. Ent.* **6**, 433-440.

Wilson, J. W., T. M. Weckwerth, J. Vivekanandan, R. M. Wakimoto, and R. W. Russell, 1994: Boundary layer clear-air radar echoes: Origin of echoes and accuracy of derived winds. *J. Atmos. Ocean. Tech.*, **11**, 1184-1206.

Wolf, W. W., J. K. Westbrook, J. R. Raulston, S. D. Pair, and P. D. Lingren, 1995: Radar observation of orientation of noctuids migrating from corn fields in the Lower Grande Valley. *Southwest. Entomol.* **S18**, 45-61

Zhang, P., S. Liu, and Q. Xu, 2005: Identifying Doppler velocity contamination caused by migrating birds part I: Feature extraction and quantification. *J. Atmos. Ocean. Tech.*, **22**, 1105-1113.

Zrnic, D.S., and A.V. Ryzhkov, 1998: Observations of insects and birds with a polarimetric. *IEEE Trans. Geosci. Remote Sens.*, **GE-36**, 661-668.