

## MESOSCALE ASPECTS IN THE FORMATION OF A PRE-CYCLOGENIC AFRICAN EASTERLY WAVE NEAR THE ETHIOPIAN HIGHLANDS

Christopher M. Hill and Yuh-Lang Lin\*  
 Department of Marine, Earth, and Atmospheric Sciences  
 North Carolina State University, Raleigh, North Carolina

### 1. Introduction

The African easterly wave (AEW) that would form the basis of Hurricane Alberto (2000) originated in the vicinity of the Ethiopian Highlands (EH) 5 days prior to cyclogenesis over the eastern Atlantic Ocean (Hill and Lin 2003, Lin et al. 2005). Initial convection began over the peaks of the northern EH during the local afternoon of 28 July 2000 (Fig. 1.1a). After dusk, the convective cells continued to grow as they tracked westward (Fig. 1.1b). By 2200 UTC 28 July 2000, growing convective cells from Eritrea moved southwestward and combined with westward-moving cells from northern Ethiopia, resulting in the development of an MCC (Fig. 1.1c). As the local nighttime progressed, the MCC peaked and began to diminish as it propagated in a southwestward and even southward direction (Fig. 1.1d). With the convection continuing to diminish, the overall cloud

feature resumes a more westward track (Fig. 1.1e). From the residual clouds, a mesovortex (MV) becomes evident over central Sudan at 1600 UTC 29 July 2000 (Fig. 1.1f). From that time, the MV would continue westward and contribute to the generation of a larger MCC over southern Chad; the National Hurricane Center first recognized the AEW by this cloud signature in a post-mortem tropical cyclone report (Beven 2000).

The development of the incipient MCC from the EH is not unlike MCC development from the Rocky Mountains (Cotton et al. 1983, Wetzell et al. 1983). Convective cells initially develop over the higher mountain peaks and are advected to a region of lower elevation with high low-level moisture, which supplies the convection for further growth. An MCC later forms over the moisture rich region during the local nighttime. The important differences between the two scenarios of




Fig. 1.1. METEOSAT-7 infrared-wavelength imagery for a) 1200 UTC 28 July 2000, b) 1600 UTC 28 July 2000, c) 2200 UTC 28 July 2000, d) 0400 UTC 29 July 2000, e) 1000 UTC 29 July 2000, and f) 1600 UTC 29 July 2000. Dot in d) depicts previous MCC center of c). Circle in f) encompasses visible MV. Imagery provided by EUMETSAT © 2003.

\* Corresponding author address: Dr. Yuh-Lang Lin, Dept. of MEAS, North Carolina State Univ., Campus Box 8208, Raleigh, NC 27695-8208. E-mail: [yl.lin@ncsu.edu](mailto:yl.lin@ncsu.edu)

MCC development are in latitude, moisture availability, and the direction of moisture advection relative to MCC propagation (e.g. Laing and Fritsch 1993, 2000). According to the Rossby radius of deformation, as given by Cotton et al. (1989):

$$\lambda_R = \frac{C_N}{(\zeta + f)^{\frac{1}{2}}(2VR^{-1} + f)^{\frac{1}{2}}},$$

where  $C_N$  is the phase speed of an inertia-gravity wave,  $\zeta$  is the vertical component of relative vorticity,  $V$  is the tangential component of wind at the radius of curvature  $R$ , an MCC from the tropics can persist if it is sufficiently large or if it exhibits sufficiently strong internal rotation, such that  $\lambda > \lambda_R$ . The development of an MV allows the incipient MCC to persist for a time, and allows for future convective development near the MV (Fritsch et al. 1994, Trier et al. 2000a, b).

Existing theories may apply for the development of the MV, observed with the pre-Alberto AEW, involving the transition from the mature stage to the dissipating stage of the MCC. Within an MCC, an MV tends to form in conjunction with the stratiform precipitation region, above where relatively high CAPE and low vertical wind shear are conducive for MV development (Fritsch et al. 1994, among others). The MV with the pre-Alberto AEW likely arises from a cyclonic PV anomaly generated between a diabatically cooled surface-based layer and a diabatically warmed upper layer (Raymond and Jiang, 1990) within the intense convection.

A 3-day average of the 850-hPa mixing ratio and wind from ECMWF operational model data, surrounding the time of observed MCC development, shows the most significant advection of low-level moisture near the west slopes of the EH from the southwest (Fig. 1.2). Not only is moisture readily available for convective development over and to the west of the EH, the direction of moisture advection promotes leftward propagation of convective cells relative to the mean-layer (northeasterly) flow (Fig. 1.3), as occurred with the northernmost and most vigorous convection observed in the presented case.




Fig. 1.2. ECMWF 3-day composite analysis of 850-hPa water vapor mixing ratio ( $\text{g kg}^{-1}$ ) and wind ( $\text{m s}^{-1}$ ) for 27 July to 29 July 2000. Contours every 2  $\text{g kg}^{-1}$ .




Fig. 1.3. ECMWF 3-day composite analysis of 600-hPa wind ( $\text{m s}^{-1}$ ) for 27 July to 29 July 2000. Contours every 5  $\text{m s}^{-1}$ .

The development of AEWs has largely been attributed to barotropic and baroclinic instabilities of the mid-tropospheric African Easterly Jet (Burpee 1972, Mass 1978, Thorncroft 1995), although AEW development has most always been described as taking place in Central or West Africa. Furthermore, the instabilities associated with the AEJ provide the means for an AEW to be sustained or to grow, and not necessarily for its generation. Carlson (1969) suggested that latent heating over significant terrain could be a mechanism for AEW generation. Being that the EH are a significant area of terrain located upstream of previously studied regions of AEW development, and being that strong convection took place over the EH for the case presented here, it stands to reason that an AEW could form near the EH.

A scarcity of observations has hindered study of AEW generation over East Africa. At best, global analyses show the synoptic patterns, and tend to miss signals of AEW generation, in this region. Weigel and Herbster (1998) visited the idea of utilizing a mesoscale model, and were apparently able to simulate an AEW from the EH region. The state of numerical weather simulation has reached the point where model output can serve as a proxy to missing observations. With this in mind, a non-hydrostatic mesoscale model is used here to simulate the environment surrounding the development of the “pre-Alberto” AEW. Primarily, the feasibility of simulating the generation of this AEW - and any other AEW - is tested.

The development of the initial mesoscale convective complex (MCC) and the associated mesovortex (MV), and the presence of an apparent lee trough, are suggested to be crucial components to the generation of the AEW. The effects of orography, sensible heating, and latent heating on the convective and kinematic development of this AEW are examined.

## 2. Methodology

The non-hydrostatic PSU-NCAR MM5 is used to simulate the environment of the area depicted in Fig. 2.1. The control simulation is conducted during the period of 00 UTC 27 July to 00 UTC 30 July 2000, during which time the incipient MCC is observed to develop, mature, and dissipate. The outer domain is set with  $253 \times 229$  grid points spaced every 12 km, and the nested domain is set with  $283 \times 283$  grid points spaced every 4 km. The two domains of the simulation are permitted to interact with one another through two-way nesting. The Kain-Fritsch cumulus parameterization (CP) scheme is used in the coarse (12-km) grid, and no CP scheme is used in the nested (4-km) grid. Vertically, each domain is structured with 45  $\sigma$ -levels. The Goddard microphysical parameterization (MP) scheme and the Blackadar boundary layer scheme are also employed. The domain boundaries are set with ECMWF 6-hourly analyses of  $1\frac{1}{2}^\circ$  grid spacing. The resolution of terrain data is 4 km for the coarse grid and 0.9 km for the nested grid.




Fig. 2.1. GLOBE Project terrain (m) within the 12-km and 4-km spaced grids. Terrain is contoured every 500 m, and shaded every 1000 m above 1000 m.

Several sensitivity experiments are to be conducted: the reduced-terrain case (RT), no PBL case (NPBL), and no latent heating case (NLH). The RT simulations utilize a terrain field with height values half of those used in the CTRL simulations, and the new initial surface fields are extrapolated downward from the former initial surface fields. The NPBL case uses no boundary layer parameterization. The NLH case excludes any CP and MP scheme.

## 3. Results

At 1200 UTC 28 July 2000, the hourly precipitation modeled by the MM5 closely aligns with the initial convection observed from satellite imagery (Fig. 3.1a; compare with Fig. 1.1a). Over time, simulated precipitation increases and accumulates westward, following the convective elements over northern Ethiopia (Fig. 3.1b; Fig. 1.1b). Notably, the convection observed over Eritrea is not captured by the MM5. Near

the time and place of the observed MCC peak, the simulated precipitation has dissipated (Fig. 3.1c; Fig. 1.1c).

As the simulated initial convection protrudes through the easterly and northeasterly flow aloft, meso- $\beta$  scale vorticity is generated near the convection and over the higher terrain of the EH (Fig. 3.2a). As the convection expands and is advected westward, the areas of vorticity similarly grow and move (Fig. 3.2b). By 2200 UTC 28 July 2000, a distinctive line of maximal vorticity along  $37^\circ\text{E}$  represents a remnant wave from the simulated convection (Fig. 3.2c).

Analysis of equivalent potential temperature ( $\theta_e$ ) from MM5 at 550 hPa (Fig. 3.3a,b) reveals that a region of low  $\theta_e$  is advected across the northern EH near the period of observed convective development. Cross-sectional analysis over Eritrea shows a surface layer of increasing  $\theta_e$  over the higher peaks and plateau of the northern EH and beneath the traversing region of low  $\theta_e$  (Fig. 3.4a); this phenomenon also occurs over northern Ethiopia. Although significant potential instability is present, simulated convection over Eritrea is weak and short-lived (Fig. 3.4b); the potentially warm low layer remains trapped near the surface, as there apparently is no simulated trigger for intense convection in this area.

The convectively generated wave simulated by MM5 can be seen along  $37^\circ\text{E}$  in a 700-hPa streamline field from the coarse domain at 2200 UTC 28 July 2000; additionally, a large cyclonic circulation is situated over southern Sudan (Fig. 3.5a). The cyclonic circulation over southern Sudan could be a lee trough; the development of this circulation may be analogous to lee vorticity development near the Central Mountain Range of Taiwan (Sun and Chern 1993, Lin et al. 1992), as well as near other topography of Africa (Mozer and Zehnder, 1996). The presence of horizontal shear also may have contributed to the dynamic development of this lee trough. By 0600 UTC 29 July 2000, the MM5 simulates the convectively generated wave combining with the lee trough (Fig. 3.5b). At this point, an AEW appears to have formed in the MM5 simulation.

The elevated heat source of the EH surface and the surrounding moisture sources of the Red Sea and the Congolese rain forests are shown to be crucial to the initiation of the scattered afternoon convection, and the eventual maturity of the convection into an MCC, with experiments of no latent heating (NLH) and no effects of the planetary boundary layer (NPBL). The reduced-terrain (RT) experiment indicates that convection alone produces a weak wave signature, and that the local orography contributes significantly to the development of the convection and of the pre-“Alberto” AEW.

## 4. Conclusions

The MM5 was able to replicate, to an extent, some but not all of the convection associated with the

developing pre-Alberto AEW. The MM5 gave a good representation of the initial convection over the peaks of EH. The fact that the MM5 misses the pivotal convective development over Eritrea could be explained by inadequate model initialization. Despite the diminutive nature of the evolving precipitation field, a traceable cyclonic wave was generated from the convection that was resolved. The simulated development of cyclonic vorticity from convection is analogous to vortical hot tower development associated with the formation of tropical cyclones (Simpson et al. 1998, Hendricks et al. 2003, Montgomery et al. 2006); for our case, cyclonic vorticity is concentrated in the middle troposphere. If the observed convection had been more completely captured by MM5, a stronger wave and an MV may have been simulated.

With satellite data being the only reliable observations, it is proposed that mesoscale numerical modeling provides the best existing means for examining mesoscale features leading to the generation of MCCs and AEWs near the EH region; in turn, satellite data best verifies the results of model simulations of the EH environment. Through a comparison between model results and with satellite data, the development of the pre-Alberto AEW appears to take place in four stages: 1) cellular convection develops in the local afternoon over the higher mountain peaks of the EH, 2) growing convective cells, and associated cyclonic vorticity, conglomerate into an MCC during the local nighttime near the lee slopes of the EH, 3) an MV develops from the remnants of the MCC, and 4) the convectively-generated MV combines with a pre-existing lee trough to form the AEW.

## 5. Future Work

More testing is needed to produce a more accurate simulation of the formation of the pre-Alberto AEW, particularly of the convective component; finer grid resolution, the inclusion of a land surface model, and land surface sensitivity testing are possible approaches to this end. Simulations of the later stages of the pre-Alberto AEW, particularly of the cyclogenesis stage, are planned.

Additional model sensitivity experiments are planned to investigate more precisely the formation mechanisms for the pre-Alberto AEW. Smoothing of the terrain data should inhibit the development of initially cellular convection, which should subsequently inhibit AEW generation. "Filling" of the Turkana Channel, or removal of the Eastern Arc Mountains of Kenya, would show the importance of any contribution of flow from the Turkana Channel to the development of the AEW. Additional cases of AEW development near the EH are also being considered for study.

**Acknowledgements.** The research contained herein has been supported by Office of Naval Research Grant #N00014-02-1-0674 and National Science Foundation Grant #ATM-0344237.

## 6. References

- Beven, J., 2000: Tropical cyclone report, Hurricane Alberto, 3-23 August 2000. National Hurricane Center, Miami, FL. (Available at: <http://www.nhc.noaa.gov/2000alberto.html> )
- Burpee, R. W., 1972: The origin and structure of easterly waves in the lower troposphere of North Africa. *J. Atmos. Sci.*, **29**, 77-90.
- Carlson, T. N., 1969: Some remarks on African disturbances and their progress over the tropical Atlantic. *Mon. Wea. Rev.*, **97**, 716-726.
- Cotton W. R., R. L. George, P. J. Wetzal, and R. L. McAnelly, 1983: A long-lived mesoscale convective complex. Part I: The mountain-generated component. *Mon. Wea. Rev.*, **111**, 1893-1918.
- \_\_\_\_\_, M.-S. Lin, R. L. McAnelly, and C. J. Tremback, 1989: A composite model of mesoscale convective complexes. *Mon. Wea. Rev.*, **117**, 765-783.
- Fritsch, J. M., J. D. Murphy, and J. S. Kain, 1994: Warm core vortex amplification over land. *J. Atmos. Sci.*, **51**, 1780-1807.
- Hendricks, E. A., M. T. Montgomery, and C. A. Davis, 2003: The role of "vortical" hot towers in the formation of tropical cyclone Diana (1984). *J. Atmos. Sci.*, **61**, 1209-1232.
- Hill, C. M., and Y.-L. Lin, 2003: Initiation of a mesoscale convective complex over the Ethiopian Highlands preceding the genesis of Hurricane Alberto (2000). *Geophys. Res. Lett.*, **30**, 36 (4 pp).
- Laing, A. G., and J. M. Fritsch, 2000: The large-scale environment of the global population of mesoscale convective complexes. *Mon. Wea. Rev.*, **128**, 2756-2776.
- \_\_\_\_\_, and \_\_\_\_\_, 1993: Mesoscale convective complexes in Africa. *Mon. Wea. Rev.*, **121**, 2254-2263.
- Lin, Y.-L., N.-H. Lin, and R. P. Weglarz, 1992: Numerical model studies of lee mesolows, mesovortices, and mesocyclones with application to the formation of Taiwan mesolows. *Meteor. Atmos. Phys.*, **49**, 43-67.
- \_\_\_\_\_, K. E. Robertson, and C. M. Hill, 2005: Origin and propagation of a disturbance associated with an African easterly wave as a precursor of Hurricane Alberto (2000). *Mon. Wea. Rev.*, **133**, 3276-3298.
- Mass, C., 1978: A numerical and observational study of African wave disturbances. Ph.D. dissertation, Univ. of Washington, 277 pp.



Montgomery, M. T., M. E. Nicholls, T. A. Cram, and A. B. Saunders, 2006: A vortical hot tower route to tropical cyclogenesis. *J. Atmos. Sci.*, **63**, 355-386.

Mozer, J. B., and J. A. Zehnder, 1996: Lee vorticity production by large-scale mountain ranges. Part II: a mechanism for the production of African waves. *J. Atmos. Sci.*, **53**, 539-549.

Raymond, D. J., and H. Jiang, 1990: A theory for long-lived convective systems. *J. Atmos. Sci.*, **47**, 3067-3077.

Simpson, J., J. B. Halverson, B. S. Ferrier, W. A. Petersen, R. H. Simpson, R. Blakeslee, and S. L. Durden, 1998: On the role of "hot towers" in tropical cyclone formation. *Meteor. Atmos. Phys.*, **67**, 15-35.

Sun, W.-Y., and J.-D. Chern, 1993: Diurnal variation of lee vortices in Taiwan and the surrounding area. *J. Atmos. Sci.*, **50**, 3404-3430.

Thorncroft, C. D., 1995: An idealized study of African easterly waves. Part III: More realistic basic states. *Quart. J. Roy. Meteor. Soc.*, **121**, 1589-1614.

Trier, S. B., C. A. Davis, and J. Tuttle, 2000: Long-lived mesoconvective vortices and their environment. Part I: Observations from the central United States during the 1998 warm season. *Mon. Wea. Rev.*, **128**, 3376-3395.

\_\_\_\_\_, \_\_\_\_\_, and W. C. Skamarock, 2000: Long-lived mesoconvective vortices and their environment. Part II: Induced thermodynamic destabilization in numerical simulations. *Mon. Wea. Rev.*, **128**, 3396-3412.

Wegiel, J. W., and C. G. Herbster, 1998: The origin of African easterly waves?. *Preprints, 16<sup>th</sup> Conference on Weather Forecasting and Analysis*. American Meteorological Society, Phoenix, AZ. 13-15.

Wetzel, P. J., W. R. Cotton, and R. L. McAnelly, 1983: A long-lived mesoscale convective complex. Part II: Evolution and structure of the mature complex. *Mon. Wea. Rev.*, **111**, 1919-1937.




Fig. 3.1. MM5 4-km grid precipitation (mm) field simulated for a one-hour period ending at a) 1200 UTC 28 July 2000, b) 1600 UTC 28 July 2000, and c) 2200 UTC 28 July 2000.




Fig. 3.2. MM5 4-km grid 600-hPa relative vorticity ( $s^{-1}$ ) and wind ( $m s^{-1}$ ) at a) 1200 UTC 28 July 2000, b) 1600 UTC 28 July 2000, and c) 2200 UTC 28 July 2000. Vorticity is shaded every  $10^{-4} s^{-1}$  up to  $4 \times 10^{-4} s^{-1}$ .




Fig. 3.3. MM5 12-km grid 550-hPa equivalent potential temperature (K) and wind ( $\text{m s}^{-1}$ ) at a) 0600 UTC 28 July 2000, and b) 1200 UTC 28 July 2000. Equivalent potential temperature is shaded every 3 K.




Fig. 3.4. MM5 4-km grid cross section of equivalent potential temperature (K) along  $14.9^\circ\text{N}$ , and between  $36.0^\circ\text{E}$  and  $40.0^\circ\text{E}$  for a) 0900 UTC 28 July 2000, and b) 1300 UTC 28 July 2000. Contours are every 1 K, and shaded every 3 K.




Fig. 3.5. MM5 12-km grid 700-hPa streamline field at a) 2200 UTC 28 July 2000, and b) 0600 UTC 29 July 2000. Dashed oval encircles signal of convective wave. Dashed line marks the apparent lee trough.