

## P2A.10 THE INTERANNUAL VARIABILITY OF TROPICAL CYCLONE ACTIVITY IN THE SOUTHERN SOUTH CHINA SEA.

Zabani Md. Zuki<sup>1</sup> George L. Limpert<sup>2</sup> Anthony R. Lupo<sup>2,\*</sup>

<sup>1</sup>Malaysian Meteorological Service  
Jalan Sultan  
46667 Petaling Jaya  
Selangor Malaysia

<sup>2</sup>Department of Soil, Environmental, and Atmospheric Science  
302 E Anheuser Busch Natural Resources Building  
University of Missouri-Columbia  
Columbia, MO 65211

### 1. INTRODUCTION

In recent years, there have been deadly tropical cyclones that have impacted the Southern South China Sea (SSCS) region, including Malaysia. Malaysia suffered direct strikes from tropical storm Greg on 25 December 1996 and tropical storm Vamei on 27 December 2001 (Chang et al. 2003). Also, the previously recorded lowest latitude for a typhoon was Sarah (3.3° N) in 1956 (Fortner, 1958). Additionally, Greg caused 238 deaths and economic losses of \$52 M in US dollars (Ooi and Ling, 1996). These storms have caused concern in Malaysia as to whether or not this was part of a long-term trend toward more frequent tropical cyclone strikes.

Tropical cyclones are among the most deadly and destructive of natural disasters in terms of loss of human life and economic destruction (Bengtsson et al., 1996). Thus, interest in research on tropical cyclones has increased substantially in recent years (e.g., Emanuel, 2005). This has been especially true regarding the study of intraseasonal and interannual variations in the intensity and occurrence of tropical cyclones, as associated with such phenomena as the Madden Julian Oscillation (MJO – e.g., Maloney and Hartmann, 2001), the Quasi-biennial Oscillation (QBO – e.g., Gray et al., 1984a,b; Chan, 1995), the El Niño and Southern Oscillation (ENSO), and the Pacific Decadal Oscillation (PDO) / North Atlantic Oscillation (NAO) (e.g., Gray et al., 1984a,b; Chan, 1985; Wu and Lau, 1992; Lander, 1994; O'Brien et al., 1996; Bove et al., 1998; Landsea et al., 1999; Pielke and Landsea, 1999; Lupo and Johnston, 2000 *hereafter LJ00*; Houghton et al., 2001).

LJ00 in their study of hurricane activity in the Atlantic region from 1944-1999 demonstrated that there was an interdecadal signal in the behavior of the ENSO response. This PDO variability could be described as there were more hurricane events per year during one phase of the PDO (e.g., 1947-1976), but little or no statistically significant ENSO variability. During the opposite phase of the PDO (1977 – 1998), there was strong ENSO-related variability in the

intensity and number of Atlantic storms.

While the most extensive studies have been performed in the Atlantic Region, there have been studies demonstrating ENSO variability in Pacific region tropical cyclone occurrence as well (e.g., Chan, 1985; Wu and Lau, 1992). These studies showed similar results for the Pacific Ocean basin as a whole. However, Ramage and Hori (1981) and Lander (1994) found no ENSO variability in the overall tropical cyclone numbers throughout the North Pacific as a whole, but significant variability in the primary tropical cyclone genesis regions. Since then, studies have shown that the focus of tropical storm activity moves southeastward (northwestward) during El Niño (La Niña) years (e.g., Chan, 1985; Wu and Lau, 1992; Lander, 1994; Wang and Chan, 2002).

Thus, this study has three objectives and they are; 1) to develop a detailed long-term climatology of tropical cyclone activity in the SSCS region, 2) to examine the character of the long-term trends and interannual variations in tropical cyclone activity, and 3) to examine variations in the background atmospheric and surface environment in the SSCS (especially in the active season – November and December). To accomplish the second objective, we examined whether or not there were significant ENSO or PDO related variations in the SSCS tropical cyclone climatology. One outcome of this study will be to produce information that may provide useful guidance for the long-range prediction of tropical cyclone occurrence for tropical weather forecasters in the SSCS region (Malaysia and adjacent nations). This will include information for advising policy makers regarding future tropical cyclone occurrences.

### 2. METHODS AND ANALYSES

#### 2.1. Analyses

The data set used here was acquired from the UNISYS website (<http://weather.usisys.com>). All tropical cyclones from 1960 – 2006 were listed on this website and included in this data set. These data were provided to UNISYS through the Joint Typhoon Warning Center (JTWC) best track data

\*Corresponding author address: Anthony R. Lupo, Department of Soil, Environmental, and Atmospheric Science, 302 E ABNR Building, University of Missouri-Columbia, Columbia, MO 65211. E-mail: [LupoA@missouri.edu](mailto:LupoA@missouri.edu).

set. The data set contains tropical cyclone categorization, name, life span, maximum wind speed, and track information. As stated in Lander (1994), tropical cyclones in the western North Pacific Ocean have been monitored by the United States Navy since 1945 and by the United States Air Force since 1947 using aircraft reconnaissance. In 1959, these entities were combined under one command, the JTWC, to track tropical cyclones over the western North Pacific. Weather satellite observations became routine in 1965, and these observations contributed to the data set as well.

The sea surface temperature (SST) information and atmospheric variables were provided by the National Center for Environmental Prediction (NCEP) and National Center for Atmospheric Research (NCAR) gridded re-analyses (Kalnay et al., 1996). These data are archived at NCAR and obtained from the National Oceanic and Atmospheric Administration (NOAA) Climate Diagnostics Center (CDC) website (<http://www.cdc.noaa.gov/cdc/reanalyses/>). These re-analyses are displayed on a  $2.5^\circ$  by  $2.5^\circ$  latitude-longitude grid available on 17 mandatory levels from 1000 to 10 hPa at 6-h intervals. These data included standard atmospheric variables geopotential height, temperature, relative humidity, vertical motion,  $u$  and  $v$  wind components, and surface information.

## 2.2. Methods

The region of study was the SCS, and it is bounded here by  $0^\circ$  to  $10^\circ$  N and  $100^\circ$  E to  $120^\circ$  E. The monthly tropical cyclone counts include those that develop locally, as well as those that enter the region from the Western North Pacific. The intensity was given as the maximum intensity attained by the tropical cyclone following the methodology of LJ00. In the Atlantic, maximum wind data have been shown to contain biases (e.g., 1944 - 1969). Landsea (1993) discusses the 1944 - 1969 bias in maximum wind speeds (as much as 5 kt or  $2.5 \text{ m s}^{-1}$ ) at some length. To reduce the influences of these biases in maximum wind speed, hurricane intensity was assigned based on the maximum Saffir-Simpson intensity attained by the storm. The statistical analysis of trends and distributions were performed using the standard statistical tests found in any standard text (e.g., Neter et al., 1988) and also follows those used in LJ00.

Many studies, such as Gray et al. (1984a), have examined the variability in the climatological background state for atmospheric quantities that have been associated with tropical cyclone development and intensification. Such variables used in other studies (e.g., Tonkin et al. 2000; Chang et al. 2005b) and included here were the area-averaged monthly; 1) SSTs and sea-level pressure (SLP), 2) the 850 – 200 hPa wind shear, 3) the low-level divergence, and 4) low-level vorticity.

The method of cycles as described by Mokhov et al. (2004) and modified by Lupo et al. (2007), was used to extract interannual and interdecadal variability from time series in this study. It is based on the analysis of simple phase plots of the first derivative of the time series versus the time series itself. The time series in this study were also filtered to extract periodicities on the interannual and interdecadal timescale, or time scales of two years or longer following the procedure in Mokhov et al. (2004). However, their publication used a two-year running mean in order to filter out the higher

frequency noise. Lupo et al. (2007) modified this technique by using a Shapiro (1970) filter.

Winds at the 850 hPa and 200 hPa geopotential height surfaces were used to generate the following quantities described below. The monthly mean vertical wind shear (850 – 200 hPa), the 850 hPa vorticity and divergence were calculated using the observed winds. Wind shear is a measure of baroclinicity in the atmosphere, and many researchers and forecasters have shown that tropical cyclone development or intensification is not favored in a strong wind shear environment (e.g. Fitzpatrick, 1997). Wind shear was used, for example, in the Statistical Hurricane Intensity Prediction Scheme (SHIPS) (e.g. DiMaria et al, 2005). SHIPS is an operational model using environmental conditions in order to forecast hurricane intensity.

The 850 hPa divergence and relative vorticity were calculated as has been done commonly in standard meteorological analysis. Also, these calculated quantities were averaged over the period of 1 November to 31 December (the active SCS tropical season), as well as area-averaged over the study region. These variables were chosen since tropical cyclone formation generally requires pre-existing low-level cyclonic disturbance for formation. These variables were also used in the study of Chang et al. (2005b) as they studied convective activity over the SCS. We used these months because such a large fraction of the tropical cyclone activity in the SCS occurred in these two months. While the time compositing was performed using the CDC website, the area averaging was carried out by downloading the data and integrating over the area using mean-value theorem (see Burkhardt and Lupo, 2005).

## 2.3 ENSO and PDO definitions

ENSO years were defined using the Japan Meteorology Agency (JMA) ENSO index to classify these as El Niño, La Niña, or neutral years. A complete description of this index can be found at the Center for Ocean and Atmospheric Prediction Studies (COAPS) (<http://www.coaps.fsu.edu>). This definition of El Niño has been used by several studies which have examined the interannual variability of atmospheric phenomenon (e.g., Bove et al., 1998; LJ00; Smith and O'Brien, 2001; Weidenmann et al, 2002), and is similar to other definitions used by other investigators (e.g., Pielke and Landsea, 1999).

Finally, the Pacific Decadal Oscillation is a 50 to 70 year basinwide variation in Pacific Region SSTs. This phenomenon is described in, for example Gershonov and Barnett (1998) in more detail, but the positive (negative) phase of the PDO represents the presence of relatively warm (cool) SSTs persisting throughout the eastern portion of the Pacific Ocean Basin. The positive (negative) phase of the PDO<sup>1</sup> persisted during the period from 1977 – 1998 (1947 to 1976), and in 1999 it was believed to have switched phases again (e.g., LJ00; Houghton et al., 2001; Lupo et al. 2007).

## 3 SCS CLIMATOLOGY

A total of 50 tropical cyclones occurred within the SCS region over the 47 year period of study (Table 1). This sample included 11 typhoons (TY, ~20%), 29 tropical storms (TS,

~60%), and ten tropical depressions (TD, ~20%). While these events constituted a very small fraction of the total Pacific Ocean basin activity (3%), two of these tropical cyclones have had recently a major impact on the nation of Malaysia. Of the 50 tropical cyclones observed over the period, 27 of these (54%) were of local origin (Table 1). Local origin here refers to storms developing within the study area described in section 2.

Table 2 demonstrated that the active tropical season for Malaysia persists from October to April. However, most of the activity (78%) occurred during the months of November (42%) and December (36%). This represented a sharp peak in the active season and likely is related to the annual migration of the Intertropical Convergence Zone (ITCZ), and the associated convection, through the SCS (e.g., Hurrell et al., 1995; Chang et al. 2005a). This seasonal peak seemed to exist regardless of whether the tropical cyclones were of local origin or they migrated into the area from the adjacent area to the east.

Tropical cyclones in the SCS region tended to move from east to west and more of these tended to move across the poleward portion of our study domain. Tropical cyclone Vamei is an example of this type of behavior. Vamei developed rapidly off the coast of Malaysia and was classified as a typhoon early on 27 December 2001 (see Chang et al. 2003). The storm then moved inland and decayed quickly over land, but not before causing widespread flooding in Malaysia and neighboring nations.

However, some storms did move west to east and tropical cyclone Greg is an example of this type of behavior. Greg developed out of a tropical depression on 24 December, 1996 and became a tropical storm just before striking northern Malaysia and then moving out of the SCS. Following the move out into the SCS, Greg was downgraded to a tropical depression again. Additionally, locally developing tropical cyclones tended also to have a short track, while most of the typhoons were storms that have migrated into the region.

Figure 1 shows a time series of tropical cyclone activity over the 47 year period. With the exception of the quiet period from 1976 – 1980, there has been steady activity in the SCS during the period. A regression analysis demonstrated that there was a slight upward trend in the number of tropical cyclones during the period of record overall (Fig. 3a). However, when this trend was tested for significance, it was found that the trend was not statistically significant at even the 90% confidence level by using the F-test and assuming that there should be no trend *a priori*. This agreed with the findings of many other studies of the active tropical regions (e.g., Free et al. 2004).

There was a prolonged period of inactivity for tropical cyclones developing within the SCS from 1974-1994 (Fig. 3b). The relatively active period since 1995 has been the result of an increase in tropical depressions developing in the SCS. The observations for the period during the 1974 - 1994 would coincide with the fact that there were more and stronger El Niño events. The observations also support the assertion of the studies cited in the reference section, mainly, that during El Niño years, the focus of tropical storm activity was located further east. Regardless, this would favor the migration of tropical cyclones into the region if there were no locally developing storms during years where one storm was observed. Alternatively, the noted increase in the

climatological trend noted above could be the result of better satellite identification techniques. However, more observations (and/or reanalysis of previous records) would be needed to confirm this hypothesis, and the satellite record may also provide better insight into trends in tropical cyclone intensity.

In order to detect interannual variations in tropical cyclone occurrences, the whole SCS dataset was partitioned into El Niño, La Niña, and neutral years. Table 3 shows that La Niña (El Niño) years were more (less) active than other years over the SCS, and this result was significant at the 95% confidence level when testing the means of these years against the mean of the entire sample using the z-score test (Neter et al., 1998) using the null hypothesis. It should be noted that since the dataset is small, statistical testing was only carried out on the complete dataset and not subsets of the data, even though the distribution of tropical storms and events of local origin were similar to the distribution of the total sample. These results, however, were similar to the results of other studies for the entire western Pacific Region as well (e.g., Wu and Lau, 1992). Finally, an examination of the variability with respect to the predominant phase of the PDO (not shown) demonstrates that there was little or no difference in the relative annual distribution of tropical cyclones. Additionally, there was no change in the interdecadal character of the ENSO related variability as found, for example, in the Atlantic by LJ00. It was not clear whether this result was a function of the small area included in the study, the small data set, or other real factors.




Figure 1. The time series of a) all tropical cyclones, and b) tropical cyclones of local origin occurring within the SCS region from 1960 – 2006. In a) the dashed line is the linear regression line.

#### 4. THE SCS BACKGROUND STATE

One goal here was to examine whether the tropical cyclone activity in active and non-active years could also be associated or correlated with variability in meteorological

parameters related to tropical cyclone formation in the climatological background of the SSCS region. In order to accomplish this, the time series of the November/December mean variables were analyzed. These included SSCS area-averaged monthly mean 1) SSTs, 2) sea level pressure (SLP), 3) 850 – 200 hPa wind shear, 4) divergence, 5) 850 hPa relative vorticity, 6) 500 hPa heights, 7) precipitable water, and 8) lifted index. The time series of variables 1 - 5 were then correlated with one another. We examined the background climatological state of the SSCS as a study of the general circulation of a much wider region is beyond the scope of this study. There are many available studies in the literature on this topic (e.g. Hurrell et al., 1995; Houghton, et al., 2001).

As expected, there were strong correlations between the SSTs and the SLP, low-level divergence, vorticity, and the wind shear (Table 4). Some of these correlations were significant at the 95% confidence level, and only the SLP correlated with the wind shear and SST correlated with vorticity showed weak correlations. Then some of these time series were filtered following the procedures in section 2 and analyzed using FFTs. A significant peak in the periodograms was found in the 3 – 7 year range which is a time period consistent with ENSO variability. A cross-spectral analysis was then performed using, for example the SST and SLP time series (not shown), and the results were consistent with the observations. Thus, it is likely that variability in SSCS background climatological state was real, and may have been due to (or at least associated with) the ENSO. Thus, the next step would be to demonstrate that during active years, the atmospheric background state must be more conducive to tropical cyclone occurrence than during non-active years. It must also be shown that the active (non-active) years were associated with La Niña (El Niño) years. Additionally, an analysis of long term trends in the SSTs and SLP (not shown) revealed that there was no statistically significant secular trend in the SSCS region.

Then, in order to test further the hypothesis (objectives – section 1, and elucidated above) and determine which primary and calculated variable(s) may have been encouraging or inhibiting tropical cyclone formation and maintenance in the SSCS, the SSTs, 200 – 800 hPa wind shear, and the 850 hPa divergence and relative vorticity were examined, and these were calculated as discussed in section 2. In order to sample these data objectively, the five most active years were chosen (Table 5) and then compared to two sets of non-active years (Tables 6 and 7 – no occurrences). The variables in Figures 2 – 4 are the mean 1 November – 31 December SLP, 500 hPa height, precipitable water, and lifted index (stability) in the SSCS region. These data were composites of the conditions within the samples shown in Tables 5-7 and these groups overall. The two sets of non-active years were compared by using the five years with the warmest SSTs and the five years with the coldest SSTs in order to generate objective samples of years with no activity and to determine the possible impact of SSTs in the SSCS. The mean SSTs for the warm (cold) non-active years was about one standard deviation (0.45° C) greater (less) than the 45 year mean (28.14° C).

The active sample contained two La Niña and three neutral years, but these three neutral years were “cold” neutral, that is SST anomalies close to La Niña conditions during the October – January period. The non-active warm







Figure 2. The mean a) sea level pressure (hPa, interval 0.1 hPa), b) 500 hPa heights (m, interval 0.5 m), c) precipitable water ( $\text{kg m}^{-2}$ , interval  $0.5 \text{ kg m}^{-2}$ ), and d) sea level pressure anomaly (hPa, interval 0.05 hPa) for November and December for the years shown in Table 5 in the SSCS region.

SST years were predominantly El Niño, but one year was a cold neutral year. The non-active cold years included one El Niño year and two “warm neutral” and one year with SST anomalies very close to zero. However, when examining all 18 years in which there was no tropical cyclone activity, 12 years were El Niño or warm neutral and two were years with SST anomalies very close to zero. The three groups described above together represent almost one third of the total number of years.

This analysis demonstrates that during the active years, the area averaged 200 – 850 hPa wind shear was of similar magnitude to that of the non-active warm SST years (Tables 5, 6) and close to the overall mean. A positive number indicated wind speeds increasing with height. The mean value for the five active years in Table 5 was smaller compared to mid-latitude values of  $15 - 30 \text{ ms}^{-1}$  found by taking several random samples of mean November/December wind shear within the mid-latitude region approximately 30 degrees latitude poleward of the study area. The wind shear values in Tables 5-7 were lower than  $15 \text{ ms}^{-1}$ , and these values are generally accepted as being supportive of tropical cyclone formation and maintenance. The relatively low tropospheric wind shear in the SSCS indicated a less baroclinic environment, especially when compared to the mid-latitudes. Additionally, an examination of the low-level relative vorticity for the active seasons demonstrated that the lower troposphere was dominated by cyclonic relative vorticity (Table 5), but the value in Table 5 was only slightly greater than the sample mean. The active year composites in Fig. 2 support the discussion above as it showed that there was relatively low SLP in the SSCS (Fig. 2a, d), and there was also ridging in the mid-troposphere (Fig. 2b), especially over the eastern portion of the SSCS. This would suggest an SSCS atmospheric environment that was more cyclonic near the surface, and the troposphere was warmer than the samples from the other group. This atmosphere would also be supportive of warm core disturbances such as tropical cyclones. Additionally, the SLP was significantly lower (at the 95% confidence level, for the whole sample  $\sigma = 0.40 \text{ hPa}$ ) than the long term mean across the SSCS, except along the equator.

However, in the non-active years with warm SSTs (Table 6), factors other than SSTs, such as the more anticyclonic environment seemed to differentiate the active-year sample from the inactive years. The SSTs from this sample were actually warmer than those for active years, but not at any accepted level of statistical significance. The area averaged low-level vorticity was the more anticyclonic (lower than the mean at the 95% confidence level) for this group (Table 6) than the two groups in Tables 5 and 7. The low-level divergence was similar to that of the active year sample. The composites of the years shown in Table 6 are given in Fig. 3. The near surface environment was clearly more anticyclonic for the non-active years (Fig. 3a), but not different significantly from the long term mean (Fig. 3). In the mid-




Figure 3. As in Fig. 2, except for the years in Table 6.

troposphere there were stronger height gradients over the region (Fig. 3b), and the flow was nearly zonal or showed weak troughing over much of the domain. This atmospheric configuration is similar to results found by Chang et al. (2005b) who found that convection was suppressed over the SSCS when the near surface environment was more anticyclonic (or as stated in their work, the ‘Borneo vortex’ was not present).

Additionally, an examination of the precipitable water suggests that the SSCS environment in these non-active years (Fig. 3c) was clearly drier than for the active year sample (Fig. 2c). Thus, it is suggested here that there was more low-level atmospheric water vapor available to potential tropical cyclones in the active years. Latent heat release has been identified as an important thermodynamic process in tropical cyclone intensification (e.g. Holland, 1997). Since climatologies of the tropics (e.g. Hurrell et al., 1995) demonstrated that most atmospheric moisture is contained in the boundary layer, the precipitable water here was taken to represent low-level moisture available for possible ingest into convective systems.

In the SSCS, it appears that even if the other background variables were similar (e.g., wind shear, low-level divergence, SSTs), the tropical cyclone activity in the region was inhibited during seasons when the relative vorticity was more anticyclonic (even though the low-level wind shear was favorable). The variables shown in Fig. 2 (Fig. 3) and Table 5 (Table 6) suggest a more (less) cyclonic and warm core background climatological state for the active (inactive) warm SST years. These results also suggest that in the SSCS, this set of El Niño years with warm SSTs may have been more hostile to tropical cyclone development as a result of the atmospheric conditions, and not necessarily because of cooler SSTs. Many studies have shown for different regions that, in addition to warm SSTs, the atmospheric background state was also a critical factor in contributing to tropical cyclone activity, (e.g., Holland, 1997; Tonkin et al, 2000; Dare and Davidson, 2004; Paterson et al. 2005). Free et al. (2004) also demonstrated that SST was not always associated with a higher potential intensity for tropical cyclones.

During the warm SST years (Table 6) in which no storms occurred in the SSCS, it is clear that the wind shear was similar to that of the active years. During the cold SST non-active years (Table 7), however, the low-level wind shear was even weaker than that of the samples in Tables 5 and 6. Nonetheless, the low-level background was also more anticyclonic (again, at the 95% confidence level) when comparing the low-level vorticities (anticyclonic relative vorticity) to those of the warm active years (Table 5). A composite map of the SLP (Fig. 4a), however, showed that in spite of weak cyclonic conditions over the ocean, similar to the active years, these values were higher than that of the long term mean (but not at the 90% confidence level). But, there were strong height gradients and troughing over the much of the SSCS region and weak ridging over the east side (Fig. 4b), which was more similar to the warm non-active years (Fig. 3b). This was also a season in which the tropospheric moisture was comparable to that of the active years (compare Fig. 4c to 2c), but more than that of the warm non-active years (Fig. 3c). Thus, in addition to the cooler SSTs, the non-active years sampled in Table 7 presented a less favorable atmospheric background environment for tropical storm




Figure 4. As in Fig. 2, except for the years in Table 7.

formation than that of the active years. Additionally, for Tables 6 and 7 (not shown), adding the next three years to these samples in which no tropical cyclone activity occurred did not change the interpretation of the results discussed here in section 4. Adding those years represented a total of 21 years sampled from the 47 year period.

## 5. SUMMARY AND CONCLUSIONS

An analysis of the interannual variations and long term trends in tropical cyclone activity from 1960 – 2006 in the SSCS was performed here in order to determine whether or not the threat of tropical cyclone activity to the nations of this region, particularly Malaysia, has increased in a detectable fashion, or whether the active period of the last decade was due to natural cycles. Malaysia was struck by two deadly tropical cyclones during the 1996 to 2001 period. There has been concern in Malaysia that tropical cyclone activity and intensity may increase in association with the expected climate change whether or not the climatic changes are naturally or anthropogenically forced. The tropical cyclone data was obtained from the UNISYS hurricane archive, and the NCEP-NCAR gridded re-analyses available via the worldwide web were used for the analysis of the climatological background conditions.

The important findings resulting from this work were;

- the SSCS experiences about one tropical cyclone event per year, whether these develop locally within the SSCS (about 54%) or they propagate into the region,
- of these, most attain tropical storm status, but relatively few become typhoons,
- the most active part of the season was confined to November and December (78% of all activity) and was likely associated with the ITCZ as it moves through the region,
- there was a slight upward trend in the occurrence of these events, but this trend was not statistically significant.

When examining the SSCS sample for interannual variability it was found that;

- La Niña (El Niño) years were more (less) active than other years, and this result was significant at the 95% confidence level when examining the total sample. The variability of tropical storms and tropical cyclones of local origin was similar to that of the total sample,
- there was no apparent climatic variability (statistically significant) in the SSCS that could be attributed to interdecadal variability such as the PDO,

A spectral analysis of the filtered climatological background variables such as SST, SLP, 200-850 hPa wind shear, 850 hPa divergence and 850 hPa vorticity showed that there was significant variability found in the 3-7 year period, which is consistent with that of the ENSO period. Examining

a subset of the most active years (all La Niña and “cold” neutral years) versus those years with no tropical cyclone activity for the five years of a) warmest SSTs (predominantly El Niño years), and b) coolest SSTs yielded some interesting results (mostly El Niño or “warm” neutral). First, during warm non-active SST years, tropical cyclone activity was likely suppressed as the low-level relative vorticity was considerably more anticyclonic, even though SSTs were about one standard deviation warmer and wind shears were similar to those of active years. The SSCS atmospheric environment for warm SST non-active years was drier than that of the active years, and did not exhibit a surface – 500 hPa structure that would be as supportive of warm-core tropical cyclones. Most of these years were also ENSO years, and two thirds of all years with no activity were El Niño or warm neutral. These results may indicate that, in the SSCS region, it was atmospheric conditions that were unfavorable to tropical cyclone development during El Niño years with warm SSTs.

During cool SST years, tropical cyclone activity was suppressed by a climatological background that was more anticyclonic at low levels than that of active years, even though the wind shear was even more favorable. The background climatological state was more similar to that of the warm SST non-active years. These results should provide forecasters in the SSCS region with guidance regarding the type of background conditions that are favorable for the occurrence of tropical cyclones. The majority of these were also El Niño and “warm” neutral years.

## 6. ACKNOWLEDGMENTS

The authors would like to acknowledge the Government of Malaysia, which provided funding for the first author. We also thank Dr. Patrick Market and Dr. Neil Fox, University of Missouri Department of Soil, Environmental, and Atmospheric Science, and Dr. Chris Wickle, University of Missouri Department of Statistics for their comments on earlier versions of this work. We’d like to thank Dr. Market for his help in drafting the figures.

## 7. REFERENCES

- Bengtsson, L., M. Boztet, and M. Esch, 1996: Will greenhouse gas-induced warming over the next 50 years lead to higher frequency and greater intensity of hurricanes? *Tellus*, **48A**, 57 – 73.
- Bove, M.C., J.B. Elsner, C.W. Landsea, N. Xufeng, and J.J. O’Brien, 1998: Effect of El Niño on U.S. landfalling hurricanes. *Bull. Amer. Meteor. Soc.*, **79**, 2477 - 2482.
- Burkhardt, J.P., and A.R. Lupo, 2005: The planetary and synoptic-scale interactions in a Southeast Pacific blocking episode using PV diagnostics. *J. Atmos. Sci.*, **62**, 1901 - 1916.
- Chan, J.C.L., 1985: Tropical cyclone activity in the Northwest Pacific in relation to the El Niño/Southern Oscillation

- phenomenon. *Mon. Wea. Rev.*, **113**, 599 – 606.
- Chan, J.C.L., 1995: Tropical cyclone activity in the western north Pacific in relation to the stratospheric Quasi Biennial Oscillation. *Mon. Wea. Rev.*, **123**, 2567 – 2571.
- Chang, C.P., P.A. Harr, and H.J. Chen, 2005b: Synoptic Disturbances over the Equatorial South China Sea and Western Maritime Continent during Boreal Winter. *Mon. Wea. Rev.*, **133**, 489–503.
- Chang, C.P., Z. Wang, J. McBride, and C.H. Liu, 2005a: Annual Cycle of Southeast Asia— Maritime Continent Rainfall and the Asymmetric Monsoon Transition. *J. Climate*, **18**, 287–301.
- Chang, C.P., C.H. Liu, and H.C. Kuo, 2003: Typhoon Vamei: An equatorial tropical cyclone formation. *Geophys. Res. Lett.*, **30**, 1150 doi10.1029/2002GL016365
- Dare, R. A., and N.E. Davidson, 2004: Characteristics of tropical cyclones in the Australian Region. *Mon. Wea. Rev.*, **132**, 3049 – 3065.
- DiMaria, M., M. Mainelli, L.K. Shay, J.A. Knaff, and J. Kaplan, 2005: Further Improvements to the Statistical Hurricane Intensity Prediction Scheme (SHIPS). *Wea. Forecasting*, **20**, 531–543.
- Emanuel, K., 2005: Increasing destructiveness of tropical cyclones over the past 30 years. *Nature*, **436**, 686
- Fitzpatrick, P.J., 1997: Understanding and Forecasting Tropical Cyclone Intensity Change with the Typhoon Intensity Prediction Scheme (TIPS). *Wea. Forecasting*, **12**, 826–846.
- Fortner, L.E., 1958: Typhoon Sarah, 1956. *Bull. Amer. Meteor. Soc.*, **39**, 633 – 639.
- Free, M., M. Bister, and K. Emanuel, 2004: Potential Intensity of Tropical Cyclones: Comparison of Results from Radiosonde and Reanalysis Data. *J. Clim.*, **8**, 1722 - 1727
- Gershunov, A., and T.P. Barnett, 1998: Interdecadal modulation of ENSO teleconnections. *Bull. Amer. Meteor. Soc.*, **79**, 2715 - 2725.
- Gray, W.M., 1968: Global view of the origin of tropical disturbances and storms. *Mon. Wea. Rev.*, **96**, 669 – 700.
- Gray, W.M., 1984a: Atlantic season hurricane frequency. Part 1: El Nino and 30 mb Quasi Biennial Oscillation influences. *Mon. Wea. Rev.*, **112**, 1649 - 1668.
- Gray, W.M., 1984b: Atlantic season hurricane frequency. Part 2: Forecasting its variability. *Mon. Wea. Rev.*, **112**, 1669 - 1683.
- \*Holland, G.J., 1997: The Maximum Potential Intensity of Tropical Cyclones. *J. Atmos. Sci.*, **54**, 2519–2541.
- Houghton, J.T., et al. (eds.), 2001: *Climate Change 2001: The Scientific Basis*. Cambridge University Press, Cambridge, UK, 857pp.
- Hurrell, J.W., H. van Loon, and D.J. Shea, 1995: Meteorology of the Southern Hemisphere. *NCAR Tech Memo no. NCAR/CAS/95-08*, Boulder, CO, 77 pp.
- Kalnay E., M. Kanamitsu, R. Kistler, W. Colins, D. Deaven, L. Gandin, M. Iredell, S. Saha, G. White, J. Wollen, Y. Zhu, A. Leetmaa, R. Reynolds, M. Chelliah, W. Ebisuzaki, W. Higgins, J. Janowiak, K.C. Mo, C. Ropelewski, J. Wang, R. Jenne, and D. Joseph, 1996: The NCEP/NCAR 40-years reanalyses project. *Bull. Am. Meteorol. Soc.*, **77**, 437-471.
- Lander, M.A., 1994: An exploratory analysis of the relationship between tropical storm formation in the western North Pacific and ENSO. *Mon. Wea. Rev.*, **122**, 636 – 651.
- Landsea, C.W., R.A. Pielke Jr., A. Mestas-Nuez, and J. Knaff, 1999: Atlantic Basin hurricanes: indicies of climate changes. *Climatic Change*, **42**, 89 - 129.
- Landsea, C.W., 1993: A climatology of intense (or Major) Atlantic hurricanes. *Mon. Wea. Rev.*, **121**, 1703 - 1713.
- Lupo, A.R., Kelsey, E.P., D.K. Weitlich, I.I. Mokhov, F.A. Akyuz, Guinan, P.E., J.E. Woolard, 2007: Interannual and interdecadal variability in the predominant Pacific Region SST anomaly patterns and their impact on a local climate. *Atmosfera*, **19**, 171 – 196.
- Lupo, A.R., and G. Johnston, 2000: The Interannual variability of Atlantic Ocean Basin Hurricane Occurrence and intensity. *National Weather Digest*, **24:1**, 1-11.
- Maloney, E.D., and D.L. Hartmann, 2001: The Madden-Julian Oscillation, barotropic dynamics, and North Pacific tropical cyclone formation. Part I: Observations. *J. Atmos. Sci.*, **58**, 2545 – 2557.
- Mokhov, I.I., D.V. Khvorostyanov, and A.V. Eliseev, 2004: Decadal and Longer-term Changes in ENSO characteristics. *I. J. Climatol.*, **24**, 401 - 414.
- Neter, J., W. Wasserman, and G.A. Whitmore, 1988: *Applied Statistics, 3rd edition*. Boston: Allyn and Bacon, 1006 pp.
- O'Brien, J.J., T.S. Richards, and A.C. Davis, 1996: The effect of El Nino on U.S. landfalling hurricanes. *Bull. Amer. Meteor. Soc.*, **77**, 773 - 774.
- Ooi, S.H., and L.K. Ling, 1996: Observational and analytical studies on tropical storm Greg affecting Sabah and its vicinity in Late December 1996. *ESCAP/WMO Typhoon Committee Annual Review 1996*.
- Pielke, R.A., and C.N. Landsea, 1999: La Nina, El Nino, and Atlantic hurricane damages in the United States. *Bull. Amer. Meteor. Soc.*, **80**, 2027 - 2033.



Ramage, C.S., and A.M. Hori, 1981: Meteorological aspects of El Nino. *Mon. Wea. Rev.*, **109**, 1827 – 1835.

Shapiro, R., 1970: Smoothing, filtering, and boundary effects. *Rev. Geophysics.*, **8**, 359 – 387.

Smith, S.R., and J.J. O'Brien, 2001: Regional Snowfall distributions associated with ENSO: Implications for seasonal forecasting. *Bull. Amer. Meteor. Soc.*, **82**, 1179 – 1191.

Tonkin, H., G.J. Holland, N. Holbrook, and A. Henderson-Sellers, 2000: An Evaluation of Thermodynamic Estimates of Climatological Maximum Potential Tropical Cyclone Intensity. *Mon. Wea. Rev.*, **128**, 746–762.

Wang, B., and J.C.L. Chan, 2002: How strong ENSO events affect tropical storm activity over the western north Pacific. *J. Climate*, **15**, 1643-1658.

Weidenmann, J.M., A.R. Lupo, I.I. Mokhov, and E. Tikhonova, 2002: The climatology of blocking anticyclones for the Northern and Southern Hemisphere: Block intensity as a Diagnostic. *J. Climate*, **15**, 3459-3473.

Wu, G., and N.C. Lau, 1992: A GCM simulation of the relationship between tropical storm formation and ENSO. *Mon. Wea. Rev.*, **120**, 958 – 977.

Table 1. Total number of tropical cyclones and annual averages over the SCS region from 1960 – 2006.

	TD	TS	TY	Total	Local Origin
total	10	29	11	50	27
Annual avg	0.21	0.62	0.23	1.06	0.57

Table 2. Total number (top row) and percentage (bottom row) of tropical cyclone occurrence in the SCS by month for the period 1960 – 2006, following the ENSO year definition used here (beginning with October).

	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Tot
T	4	21	18	1	1	1	2	2	50
%	8	42	36	2	2	2	4	4	100

Table 3. The average annual tropical cyclone activity separated by a) El Niño years for all tropical cyclones (All), typhoons (TY), tropical storms, tropical depressions (TD), and tropical cyclones of local origin.

	All	TY	TS	TD	Local Origin
Neutral	1.1	0.3	0.6	0.3	0.6
El Niño	0.4	0.2	0.2	0	0.3
La Niña	1.6	0.1	1.2	0.3	1.0
All Years	1.0	0.2	0.6	0.2	0.6

Table 4. The correlation between SSTs, SLP, geostrophic wind shear, wind divergence, and vorticity. The asterisk denotes a correlation at a 95% confidence interval.

Parameter	SST-SLP	SST-Div	SST-Shear	SLP-Div	SLP-Shear	Div-Shear
Correlation	-0.59*	-0.70*	0.29	0.55*	-0.10	-0.23

Parameter	SST-vor	Div-Vor	Shear – Vor
Correlation	0.15	-0.53*	0.35

Table 5. The area-averaged November and December means for the five most active tropical seasons in the SCS.

Year	Total TCs	ENSO Year	SSTs (°C)	wind shear (ms <sup>-1</sup> )	divergence (x 10 <sup>-6</sup> s <sup>-1</sup> )	relative vorticity (x 10 <sup>-6</sup> s <sup>-1</sup> )
1996	4	Neutral	28.6	11.4	-2.26	5.70
1998	3	La Nina	28.1	12.4	-2.49	10.00
1970	3	La Nina	28.1	12.0	-2.22	-2.35
1962	3	Neutral	27.7	9.8	-0.03	-2.10
1995	2	Neutral	28.3	9.4	-2.44	5.85
		means	28.1	10.1	-1.88	3.42

Table 6. As in Table 5, except for five non-active years with the warmest SSTs.

Year	Total TCs	ENSO Year	SSTs (°C)	wind shear (ms <sup>-1</sup> )	divergence (x 10 <sup>-6</sup> s <sup>-1</sup> )	relative vorticity (x 10 <sup>-6</sup> s <sup>-1</sup> )
1987	0	El Niño	29.0	9.2	-1.95	0.90
2002	0	El Niño	28.8	2.7	-1.31	-0.90
1966	0	Neutral	28.4	12.8	-2.62	-2.25
1963	0	El Niño	28.4	7.5	-0.08	-4.95
1986	0	El Niño	28.3	5.8	-2.94	1.55
		Means	28.6	8.5	-1.78	-1.13

Table 7. As in Table 6, except for five non-active years with the coldest SSTs.

Year	Total TCs	ENSO Year	SSTs (°C)	wind shear (ms <sup>-1</sup> )	divergence (x 10 <sup>-6</sup> s <sup>-1</sup> )	relative vorticity (x 10 <sup>-6</sup> s <sup>-1</sup> )
1984	0	Neutral	27.3	4.9	-1.44	-0.75
1978	0	Neutral	27.5	6.2	-0.35	0.16
1979	0	Neutral	27.9	7.1	0.37	-0.49
1977	0	Neutral	27.9	4.8	0.16	-1.40
1991	0	El Niño	27.9	2.9	-0.02	-0.68
		Means	27.7	5.2	-0.26	-0.63