

## DO CITIES ENCOURAGE WARM-SEASON THUNDERSTORM FORMATION AND INTENSIFICATION?

Walker S. Ashley<sup>\*,\*</sup>, Mace L. Bentley<sup>\*,</sup>, and Tony Stallins<sup>#</sup>

<sup>\*</sup>Meteorology Program, Department of Geography, Northern Illinois University, DeKalb, Illinois

<sup>#</sup>Department of Geography, Florida State University, Tallahassee, Florida

### 1. INTRODUCTION

Anthropogenic land use/land cover (LULC) changes modulate meteorological and/or climate forcings at the local, regional, and global scales. The IPCC, NRC, and U.S. Climate Change Science Program have emphasized the importance of LULC changes on influencing human-environment systems. The most dramatic LULC modification of the natural environment is arguably urban development. Currently, over 50% (82%) of the world (U.S.) population lives in an urban area, and this proportion is predicted to grow beyond 70% (90%) by 2050 (UNFPA 2007).

There is a large body of literature that illustrates that urban areas modify their weather and climate (Shepherd 2005). With the advent and improvement of remote sensing platforms over the last two decades, thunderstorm climatologies of major metropolitan areas have been developed through the employment of cloud-to-ground lightning data, space-borne precipitation radar data, and, recently, National Weather Service WSR-88D radar data. However, no long-term, multi-scaled investigations of urban-enhanced convective storms exist. In addition, none of these prior studies incorporate "controls," regions to reveal whether urban convective anomalies are reproduced in rural regions. A major question in urban weather and climate research is how much cities vary in their propensity to modify local convection.

Our research illustrates how urbanization augments warm-season thunderstorm activity among a range of cities in the southeastern U.S., a region that experiences diurnally forced convection during the warm-season with only occasional synoptic-scale forcing for ascent. Urbanization contributes to a variety of processes that may lead to enhancement of thunderstorms; these mechanisms include urban heat island (UHI) dynamics, enhanced localized forcing due to increased surface roughness, and elevated production of aerosols and, subsequent, cloud condensation nuclei. By comparing the timing, location, and intensity of convection in cities and adjacent control regions, we analyze how these demographic and LULC changes feed back to strong and severe thunderstorms.

### 2. DATA AND METHODOLOGY

The seven locations in the southeastern U.S. chosen for investigation comprise five cities of various sizes with the addition of two control, or null, regions that are predominantly rural (Table 1). We limited our analyses to the warm-season months of June through August; the part of the year when the Southeast U.S. is largely absent synoptic-scale forcing and is dominated by a maritime tropical airmass associated with weakly forced, diurnally driven convection. We hypothesize that the formation of isolated, airmass storms in these conditions across our domain will be coupled to LULC, including urban land use effects. Only city-specific days where the spatial synoptic classification indicated a moist tropical (MT), moist tropical plus (MT+), or moist tropical plus-plus (MT++) airmass were examined (Table 1). Because our rural control locations were not available in the SSC database, we used the closest SSC identified city.

In order to examine urban effects on thunderstorm frequency and magnitude across these locations, we assessed 10 years (1997-2006) of NOWrad national composites of WSR-88D reflectivity data produced by WSI Corp. These data have temporal, spatial, and reflectivity resolutions of 5 minute, 2km x 2km, and 5dBZ, respectively, and illustrate the largest reflectivity measured in a 5-minute interval by any radar scan in a column above each grid cell. The data are quality controlled by WSI to filter artifacts such as ground clutter. In order to assess the intensity of thunderstorms, we examined the spatiotemporal constituents of three reflectivity categories: 1) the amount of time that a grid cell experienced 40, 45, or 50 dBZ, which illustrates the approximate reflectivity magnitudes of a modest, nonsevere thunderstorm, 2) the amount of time that a grid cell experienced greater than 50 dBZ, which is associated with strong-to-severe thunderstorms, and 3) the "sum" total of the medium and high reflectivity events, or the total amount of time that a grid cell experienced greater than 39 dBZ. We supplemented our radar analyses for the Atlanta, GA MSA with 10 years (1997-2006) of cloud-to-ground lightning data acquired from the U.S. National Lightning Detection Network (NLDN) produced by Vaisala Corp.

To examine a variety of definitions of urban and rural areas, and their influences on our results, we used three separate methods to discriminate between these fundamental LULC types on the aforementioned 2km x 2 km base grid across the domains (Figure 1).

---

\* Corresponding author address: Dr. Walker Ashley, #118 Davis Hall, Meteorology Program, Dept. of Geography, Northern Illinois University, DeKalb, IL 60115; email: washley@niu.edu; web: chubasco.niu.edu

First, we employed the National Land Cover Database (USGS 2001a; hereafter, NLCD), which contains standardized land cover components using Landsat 5 and 7 data. In particular, we identified the urban region of our largest cities (Atlanta, Memphis, and Birmingham) that contained four of NLCD's land cover classifications, including: Developed, Open Space (21), Developed, Low Intensity (22), Developed, Medium Intensity (23), and Developed, High Intensity (24). Each 2km x 2km grid cell that contained an areal coverage of these four NLCD classifications greater than 49% was considered "urban"; cells that contained less than or equal to this critical threshold were considered "rural". Those rural areas contained within the urban corridor demarcation (i.e., rural grid cells enclosed in the larger urban polygon) were considered urban due to their containment in the larger urban core outline.

Next, we used the USGS's (2001b; hereafter, USGS) urban areas vector layer dataset, which illustrates areas of built-up land where large populations exist. The urban corridor was assessed in the same manner as above; essentially, if an individual grid cell contained greater than 49% of this vector classification, it was considered urban.

Finally, we employed a set of simple geometric buffer constructions based on spatial satellite examination of recent development trends to illustrate the urban corridor for all cities examined, as well as the two rural controls (Table 1). The control regions had surrogate urban areas defined by a 31km buffer, which was the average radius of the five urban buffers used for the cities.

**Table 1.** Characteristics of MSAs and control regions examined, ranked by population (U.S. Census). Buffers in km.

MSA or control	Population (2009 est.)	1990-2009 % $\Delta$	Urban Buffer	Rural Buffer	MT Days Examined
Atlanta, GA	5,475,213	84.98%	40	100	362
Memphis, TN	1,304,926	29.55%	35	87.5	471
Birmingham, AL	1,131,070	34.63%	30	75	429
Jackson, MS	540,866	36.79%	25	62.5	427
Montgomery, AL	366,401	25.26%	25	62.5	441
Starkville, MS control	-	-	31	77.5	438
Heflin, AL control	-	-	31	77.5	429

### 3. RESULTS

#### 3.1 Spatial Analysis


Visual analysis of the warm-season, maritime tropical convective reflectivity characteristics across cities and controls revealed a trichotomy in convective enhancement due to urban influences (Figure 1 and 2). The largest MSAs – Atlanta, Birmingham, and Memphis – had medium, high, and total (medium + high) reflectivity occurrence (i.e., number of 5-minute scans that meet the dBZ criterion ) and day (i.e., number of

days that meet the dBZ criterion) hotspots clustered within and near the urban corridors, while analyses of the smaller cities (Jackson and Montgomery) suggest a more indistinct urban influence. The control regions (Heflin and Starkville) illustrate reflectivity trends that have little to no spatial correlation with surrogate city centers. This initial spatial interpretation suggests that: 1) large cities do instigate thunderstorm formation as well as augment convective intensity; and 2) the scale and distribution of the city footprint is an important factor in determining urban influences on thunderstorm formation and sustenance. Analyses of radar reflectivity trends suggest that cities must be of sufficient size to identifiably increase thunderstorm frequencies and that, ultimately, there appears to be a tipping point at which humid subtropical cities and their heating nodes, roughness factors, and/or aerosol production become significant in augmenting convection.


Atlanta and Birmingham's reflectivity hotspots have different spatial densities that appear to be controlled by the distribution of population and development across the two urban landscapes. Birmingham has a more concentrated urban core in comparison to Atlanta's well-known sprawling quality. These characteristics appear to be manifested in the thunderstorms amplified by the cities. Birmingham has a much more concentrated occurrence and day reflectivity hotspot, with Atlanta revealing a broad elevated distribution of enhanced reflectivity across the urban core and surrounding suburbia. Birmingham's unique orography has shaped both urban and suburban development trends and appears to influence the area's thunderstorm distribution. While it is possible that orography could modulate the urban-enhancement found in Birmingham, it is hypothesized to have a geographically contingent effect in comparison to urban LULC.

Converse to Atlanta's sprawl development mode, Memphis' population expansion and, therefore, urban thunderstorm influences are sharply confined to the urban core with a broad decreasing enhancement that slopes eastward away from the CBD. This enhancement pattern parallels the underlying urbanization and suburbanization trends found in this MSA, which are largely confined to the city center and eastward due to the Mississippi River and associated agriculturally rich flood plain encountered directly west of the CBD.

Spatial reflectivity characteristics of the Jackson and Montgomery regions demonstrate complex local and/or regional controls on thunderstorms. Analysis reveals that the urban area itself appears large enough to exhibit some influence on thunderstorm patterns -- evidenced especially in Montgomery's total reflectivity days, but also more subtly in Jackson's few high-activity pixels within and near the city center. However, the urban enhanced dynamics appear not sufficient enough to overcome other local/regional processes (e.g., LULC effects, ambient atmospheric environment characteristics at initiation, etc.) to make urbanization a



**Figure 1.** Frequency of June-August medium and high reflectivity days, or total days  $\geq 40$  dBZ in each 2-km grid cell, for the 10-year period of record for a) Atlanta, b) Memphis, c) Birmingham, d) Jackson, e) Montgomery, f) Starkville control, and g) Heflin control. Red circles correspond to urban-rural buffers outlined in Table 1. For a-c, the blue (green) lines represent NLCD (USGS) urban delineations.



**Figure 2.** As in Fig. 1, except frequency of June-August medium and high reflectivity occurrences, or total number of 5-minute occurrences  $\geq 40$  dBZ in each 2-km grid cell, for the 10-year period of record.

dominant influence on thunderstorm formation and sustenance. This leads to a noisier solution in the medium city climatologies as individual thunderstorm occurrences and days likely contain an urban signature combined with, and possibly overshadowed by, other localized forcing mechanisms modulating the convection.

Rural controls -- in our investigation, Heflin and Starkville -- permit the validation of the extent to which convective anomalies witnessed in proximity to urban centers are reproduced in rural regions, whether by natural variability or other environmental features such as topography. Visually, the reflectivity patterns found in the Heflin, which straddles Atlanta and Birmingham MSAs, reveal no singular hotspots centered near the surrogate urban core. Starkville has a broad maximum to the north as well as a spine of higher reflectivity that appears tied specifically to non-urban LULC effects. In particular, the narrow north-south correspondence of the physiographic-distinct land covers of Upper Coastal Plain, Blacklands, and Interior Flatwoods in the Starkville region appear to be the dominating influence on the warm-season convective climatology. Further analysis of this region's physiographic influence on convection will be required to determine how nonclassical mesoscale circulations induced by unique land-cover-type boundaries may be augmenting the convective climatology. Ultimately, initial conclusions from the mid-sized city spatial analysis is a reminder that urbanization is only a single, albeit increasingly important, variable in determining the complex distribution of thunderstorms.

### **3.2 Statistical comparisons**

To objectively assess the initial findings revealed in the spatial analysis, we examined the statistical significance for differences in the mean (median) reflectivity characteristics using the independent-samples t-test (Mann-Whitney U Test) between the rural and urban regions across the seven domains. Significance testing ( $\alpha = 0.05$ ) across various medium, high, and sum reflectivity categories revealed that, in most instances, large cities had statistically significant increases in warm-season thunderstorm occurrences and days in comparison to their rural counterparts. We examined this significance using the three separate methods for defining urban and rural areas as discussed in Section 2.

All three of the largest cities examined (Atlanta, Birmingham, and Memphis) had statistically significant increases across all urban delineations except for a single case -- medium occurrences in the Birmingham area using the buffer method, which revealed a significant decrease in medium reflectivity occurrences ( $t=-3.49$ ;  $p=0.000$ ). Further analysis of the spatial distribution of medium occurrences revealed that this outlier may be caused by two reasons: 1) the urban buffer demarcation of 30km from city center may be too generous for this more focused population center, and 2) an area of enhanced medium reflectivity occurrences

found in the southeast quadrant of the rural domain tended to increase the rural reflectivity contribution. It may also be that the particular nature of aerosols in Birmingham -- specifically, their size and concentration -- may result in a different distribution of medium and high reflectivity occurrences. This significant decrease in the Birmingham medium occurrence data was not reflected in the high occurrences, but did manifest when medium and high occurrence data were summed. In this particular sample, there was not enough evidence to conclude that the mean rural and urban event occurrences  $\geq 40$  dBZ were statistically distinguishable.

The smaller cities of Jackson and Montgomery had a range of results. Across most reflectivity categories, Jackson rural and urban buffers had no significant differences in the means or medians, which confirms the initial hypotheses that the Jackson urban area is perhaps too small to modify thunderstorms or is masked by other nonclassical mesoscale circulations induced by non-urban LULC effects. While Jackson is relatively close (250km) to the Gulf of Mexico, there appears to be little evidence that the sea-breeze effect extends this far inland recurrently and, therefore, is influencing the climatology found herein.

Montgomery has statistically greater reflectivity counts in its 25 km urban buffer in comparison to the surrounding rural region except for the category of medium days ( $t=0.74$ ,  $p=0.460$ ). This statistical evidence reconfirms the apparent moderate urban influence identified visually, suggesting that despite the cities relatively small size, it may impact warm-season thunderstorm frequency.


Heflin and Starkville have differing statistical results, which must be weighed against visual analyses of the reflectivity climatology to confirm the lack of urban influence. In Heflin's case, statistical tests suggest that the frequency of various reflectivity characteristics for the surrogate urban area and rural region were not statistically distinguishable or had significantly lower "urban" reflectivity in comparison to the rural region. This result implies that the Heflin region exemplifies no urban influence, providing support to the hypothesis that urbanization influences the thunderstorm climatology in and around cities of sufficient size. Across all various reflectivity characteristics, Starkville's surrogate urban sphere was significantly higher compared to its rural counterpart. This result would initially appear to be counter to the hypothesis that urbanization does augment thunderstorm frequencies and intensity. A visual inspection of the reflectivity climatologies suggests that the differences found are not linked to urbanization effects since urbanization is minimally developed. Locally variable physiography may, in this case, have produced the statistical anomaly.

### **3.3 Graphical analyses of urban convection**

Employing a set of scatterplots and quadratic line fits further elucidates the urban influence on thunderstorm occurrences across the domains. In particular, we

examined the frequency of various reflectivity thresholds, as well as lightning strikes for Atlanta, versus distance from city center to appraise the urban effects on thunderstorms. A concave curvilinear form sloping away from a city center supports the hypothesis of urban enhanced thunderstorms in and near city center.


The resulting medium and high, or total, reflectivity scatterplots and line fits confirm the urban effects on thunderstorm frequency evident in our other analyses (Figure 3). We present only the total reflectivity scatterplots of Atlanta using the NLCD demarcation, but provide comprehensive graphical summary of the line fits for all cities and null regions in Figure 4. Both Atlanta occurrence and day scatterplots exemplify frequency counts that slope dramatically away from the high values near the CBD to the comparatively low values found across the rural area. The Atlanta buffer delineation of 40km, as well as NLCD-based urban definition, appear to separate well the urban and rural frequency shifts since the slopes of the quadratic fits appear to lessen near the urban/rural separation.



**Figure 3.** Quadratic line fits and associated scatterplots representing the frequency of a) total days  $\geq 40$  dBZ and b) total number of 5-minute occurrences  $\geq 40$  dBZ for each 2-km grid cell versus distance from city center in the Atlanta domain for the 10-year, June-August period of record. NLCD urban delineated cells are colored red, whereas non-urban cells are blue. The 40-km urban buffer for Atlanta is represented as solid line emanating from the x axis.

Atlanta and Memphis slopes are some of the steepest found across the domains, which implies that the urban effects on convection in these cities are the most dramatic. Birmingham continues this large-city tendency for total days, but is inconclusive for occurrences. Again, this gradual slope is due to the enhanced counts found in southeast Birmingham domain for medium occurrences, which is reflected as a flat line fit (not shown) for this classification. Individually, Birmingham's medium days, high days, and high occurrence data (not shown) reveal steep slopes of elevated reflectivity counts near the city center, and reduced counts across the rural domain.

The mid sized cities Montgomery and Jackson continue to exhibit mixed results in terms of slope magnitude, but do illustrate a broad decrease in reflectivity counts away from the city centers. Montgomery slopes for medium plus high days and events are more substantial; in fact, the Montgomery total occurrences is the steepest slope of all domains. Both Starkville and Heflin curves are either flat or convex, which illustrate a different distribution when compared to the city domains. Despite Starkville's decreasing slope from "city" center, its diminishing trend and lack of substantial concavity does not fit the pattern found in the urban cases. Heflin is uniquely flat for reflectivity occurrences, and slope ascends from the surrogate city center for day counts, an atypical characteristic that supports the lack of urban influence near the domain's center.



**Figure 4.** Quadratic line fits representing the relative frequency of a) total days  $\geq 40$  dBZ and b) total number of 5-minute occurrences  $\geq 40$  dBZ versus distance from city center for the seven domains examined. Various urban-rural buffers used (see Table 1) are represented by black line segments.



### 3.4 Percentage increase in thunderstorms

Comparing the means of the urban and rural reflectivity frequencies illustrates the proportional increases, or decreases, in thunderstorms across domains. Atlanta and Memphis featured percentage increases in medium, high, and total reflectivity frequencies across the urban areas. In both cities, the largest percentage increases ( $\geq 36\%$  positive augmentations) were associated with high-end occurrences and days, suggesting that these urban areas are not only leading to an increase in thunderstorm frequency, but elevated occurrences of intense thunderstorm events.

**Table 2.** Percentage change in the mean frequency of reflectivity characteristics examined for different various urban delineation methods. Percentage changes in lightning counts and days are represented for Atlanta.

City (urban delineation)	Percentage Change in Reflectivity						Lightning	
	Med Occur.	High Occur.	Total Occur.	Med Days	High Days	Total Days	Flash Count	Flash Days
Atlanta (NLCD)	17.7	42.9	18.6	9.0	36.4	11.7	37.7	19.5
Atlanta (USGS)	23.7	61.9	25.0	14.0	54.5	17.0	41.7	18.1
Atlanta (Buffer)	16.7	50.0	17.8	8.0	36.4	10.8	34.2	13.6
Memphis (NLCD)	25.0	47.1	25.9	20.2	55.6	23.3	-	-
Memphis (USGS)	20.7	47.1	21.5	22.3	40.0	24.0	-	-
Memphis (Buffer)	7.5	43.8	8.6	8.6	33.3	10.7	-	-
Birmingham (NLCD)	3.3	55.9	5.5	11.6	33.3	14.3	-	-
Birmingham (USGS)	7.0	70.6	9.6	15.5	44.4	19.0	-	-
Birmingham (Buffer)	-1.5	20.6	-0.6	6.3	11.1	7.5	-	-
Jackson (Buffer)	-0.6	0.0	-0.6	2.0	0.0	0.9	-	-
Montgomery (Buffer)	2.6	16.7	3.1	0.7	15.8	2.5	-	-
Starkville (Buffer)	11.1	32.0	11.8	4.0	28.6	6.5	-	-
Heflin (Buffer)	0.8	-17.2	0.5	-1.5	-12.5	-2.7	-	-

Birmingham demonstrates similar percentage increases, with high-reflectivity occurrences and days ranging from 11-70% increases in the urban versus rural regions. As discussed prior, the medium occurrences for Birmingham are conflicting, suggesting negligible change. Whether this is simply a reflection of the enhancement of moderate thunderstorm activity due to Gulf of Mexico proximity, or some other effect due to LULC-induced mesoscale circulations (e.g., effects of nearby Mitchell and Lay Lakes and associated lake breezes) is difficult to discern. Summing medium and high reflectivity days suggests that Birmingham experiences 7-19% more thunderstorm days in comparison to its rural surroundings; this is comparable to the urban domain percentage increases found in Atlanta (11-17%) and Memphis (11-24%) for total days.

Jackson experiences minor percentage changes from urban to rural; whereas, Montgomery features neutral to positive change, with high days and occurrences having the greatest differences (~16% positive augmentation). The null cases of Starkville and Heflin have conflicting results, with Heflin experiencing negative to neutral thunderstorm percentages changes across the surrogate urban center and Starkville experiencing net percentage increases across the urban proxy. The mid size city and control city results are expected considering the prior conclusions drawn from the spatial and statistical perspectives.

Finally, examining 10 years of lightning data for the Atlanta domain supplements the strong evidence of urban modification of warm-season thunderstorm frequency and magnitude. Using the three urban-delineation methods, the Atlanta urban area had a 34-42% increase in the mean lightning flashes per grid cell in comparison to the surrounding rural region. Lightning was, on average, 14-20% more common on any given day in the urban region versus rural region.

## 4. CONCLUSION

Past research has demonstrated that large cities can augment sensible weather effects due to various urban-induced kinematic, thermodynamic, and pollution effects (Shepherd 2005). This prior body of work has focused principally on temperature and precipitation increases due to the aforementioned effects. Our study builds upon these efforts by presenting a high-resolution, radar-based climatology of warm-season convection for a multi-city region in the southeastern U.S., an area that has experienced considerable increases in urban and suburban populations over the last few decades. We characterized the frequency of moderate and high-end thunderstorms in an effort to see how cities reshape the distribution of convective hazards. Through this quantitative effort, we identified the scalar extents related to urban-enhanced convection. We implemented experimental controls, which has been cited as a major shortcoming in historical studies of urban-induced precipitation (Shepherd 2005).

Specifically, our results illustrate substantive evidence of urban effects on thunderstorm frequency and severity for major cities as well as evidence of non-urban LULC effects at a control site. The mid-sized cities of Jackson and Montgomery had more subtle urban effects, suggesting that urban impacts on convection are more muted and, therefore, the urban-enhanced dynamics are not significant enough to dominate other local/regional processes -- i.e., the urban effects are embedded among larger controls. Therefore, the extent and shape of urban land cover can be seen as positively correlated with the potential for urban-augmented thunderstorms.

As cities continue to grow into the 21st Century (UNFPA 2007), so will the convective feedbacks and, in return, enhanced thunderstorm risk they engender. When this

risk is juxtaposed with elevated physical vulnerability created by urban infrastructure (e.g., impervious surfaces, outdated and aging storm drainage infrastructure, etc.), as well as the social vulnerability due to a concentration of millions of people and their assets into these centers, devastating consequences can result. The 2009-10 flooding episodes in Atlanta, Nashville, and Oklahoma City are illustrative of the disastrous consequences that can occur when heightened risk and vulnerability commingle at the scale of a metropolitan region.

## **5. ACKNOWLEDGEMENTS**

The authors wish to thank Phil Young and Rick Schwantes of the NIU Advanced Geospatial Laboratory for their technical expertise and equipment. Portions of this research were supported by National Science Foundation Grant #0649343.

## **6. REFERENCES**

Shepherd, J.M., 2005: A review of current investigations of urban-induced rainfall and recommendations for the future. *Earth Interactions*, **9**, 1–27

UNFPA, 2007: State of the World Population: Unleashing the Potential of Urban Growth United Nations Population Fund <http://www.unfpa.org/swp/2007/>

U.S. Geological Survey, 2001a: National land cover data set 2001. Reston, Virginia.  
<http://seamless.usgs.gov/products/nlcd01.php>

U.S. Geological Survey, 2001b: Urban Areas of the United States. Reston, Virginia.  
<http://nationalatlas.gov/atlasftp.htmlr>