

THE ELECTROMAGNETIC-HYDRODYNAMIC ANALOGY: AN APPROACH TO VORTEX DYNAMICS AND PRESERVATION IN TORNADIC SIMULATIONS

Marcus L. B ker *
Western Illinois University, Macomb, IL

Gregory J. Tripoli
University of Wisconsin-Madison

1. INTRODUCTION

Theoretical development regarding vortex dynamics within atmospheric turbulence has been one of the most challenging scientific problems over the last century. Given the breadth of application, it is clear why there is such an interest in the subject. However, while incremental advances have been made in recent years (with the help of exponentially improving computational resources) there has not been a clear paradigm shift in some time. Even so, there is a slowly growing body of literature utilizing the potentially synergistic relationship between the fields of electro-magnetism (EM) and hydrodynamics (HD). This synergy is based on the similarity between the equations that govern fluid dynamics and the equations that govern electro-magnetism. Thomson (1931) was one of the first to employ this type of ‘‘analogous thinking’’ when developing a visual framework for electromagnetism. A table of analogous variables has been compiled from several sources (Belevich, 2008; Marmanis, 1998, Pinheiro, 2009) in Fig. 1. Exploring and exploiting this analogous relationship may be a catalyst for a radical shift in how coherent structure in fluids is studied.

2. VORTEX DYNAMICS: METHOD AND DIAGNOSTICS

One of the main struggles in the numerical modeling of tornadoes is the tendency for over-production of turbulent diffusion in the vicinity of the vortex. Several methods have been developed to combat this numerical loss of kinetic energy, including ‘‘vortex confinement’’, (e.g. Steinhoff and Underhill, 1994) and other antidiffusion methods. However, it seems desirable to find physical reasoning for why a tornado does not break down into turbulence, given the extreme deformation fields near the vortex. Observations show that there can be coherent organization, subsequent breakdown, and reorganization of the parent vortex and surrounding vortex filaments (e.g. Rotunno, 1984).

In order for vortex filaments to merge into the parent vortex, they must (1) be aligned with the parent vortex and (2) be advected toward (or into) the core of the parent vortex. Using the Lamb vector and some of the

relationships well-known in electromagnetism, we examined the behavior and self-organization potential of vortex filaments surrounding a tornado in a high-resolution simulation.

We use the University of Wisconsin Nonhydrostatic Modeling System (UWNMS: see Tripoli, 1992), with up to 5 nested grids, obtaining horizontal and vertical resolution below 25 meters. An idealized vertical sounding, based on a tornadic supercell environment, is used to initialize the meteorological fields.

One can imagine randomly oriented small-scale vortices (or angular moments) embedded within the larger scale rotation of the tornado. By decomposing the Lamb vector into mean and perturbation components, we find only one combination (using the mean vorticity and perturbation velocity) will geometrically yield a torque across a segment of a perturbation vortex filament embedded within a large-scale vorticity field

$$\frac{\partial \mathbf{u}'}{\partial t} \approx -(\bar{\boldsymbol{\omega}} \times \mathbf{u}') \equiv -\mathbf{I}'_a \quad (1a)$$

where \mathbf{u}' is the perturbation velocity, $\bar{\boldsymbol{\omega}}$ is the mean vorticity, and \mathbf{I}'_a is this component of the Lamb vector.

The magnitude of the curl of this term, $|\nabla \times \mathbf{I}'_a|$, is used to diagnose ‘‘gyroscopic alignment torque’’, which is the HD analog to the torque on a magnetic dipole moment immersed in a large-scale magnetic induction:

$$\boldsymbol{\tau} = \boldsymbol{\mu} \times \mathbf{B} \quad (1b)$$

where \mathbf{B} is the large-scale magnetic induction, and $\boldsymbol{\mu}$ is the magnetic moment of the small dipole. These parameters are illustrated in Fig. 2.

Regarding the alignment of vortex loops, Stokes Theorem states that vortex lines cannot just ‘end’ in the middle of a fluid; they must either ‘loop back on themselves, or terminate at a surface. That does not preclude alignment of vortex segments, however. One can envision a poloidal configuration of vortex lines aligning themselves within the tornado.

We investigate another EM-analogous parameter which is related to a mechanism in terms of hurricane prediction (known as the ‘‘beta effect’’). We term this new

* Corresponding author address: Marcus L. B ker, Western Illinois University, Geography Department, Macomb, IL, 61455; e-mail: ml-buker@wiu.edu.

parameter (which is a *vector*) the “vortex beta force”, β . Hurricanes (small vortices) are known to be drawn upgradient toward higher same-signed vorticity (as the Coriolis parameter increases with higher latitude) through the conventional “beta-effect”. This is absolutely consistent with this new β parameter, and it is directly analogous to the same mechanism responsible for “magnetic shielding” of charged particles: the force on the particle is proportional to the gradient of the configuration of vorticity aligned with the particle’s magnetic moment:

$$\mathbf{F} = \nabla(\boldsymbol{\mu} \cdot \mathbf{B}) \quad (2a)$$

Substituting in the analogous variables, one obtains a formula for the force on a small local moment, which is generally aligned with the large-scale rotation:

$$\mathbf{F} = \nabla(\boldsymbol{\mu} \cdot \bar{\boldsymbol{\omega}}) = \beta \quad (2b)$$

We investigate these two analogous formulations of vortex interaction and discuss them in the following section.

3. “EM-LIKE” TORQUING AND MERGING: RESULTS AND DISCUSSION

3.1 Gyroscopic torque

To simplify the gyroscopic torque formulation, we looked at the mathematical formulation for the electromagnetic torque, and applied this analogously to the vortex interaction problem. To calculate this new parameter, information was needed about the **LOCALLY** defined angular momentum. Since the pressure gradient force is the only force (besides friction) that accelerates the flow in a Lagrangian sense, the normal component of pressure gradient acceleration to velocity was computed, and from this a local radius of curvature was obtained. Obviously, there is a strong correlation of local angular momentum and the local curvature radius:

$$\boldsymbol{\mu} = \mathbf{r} \times \rho \mathbf{u} \quad (3)$$

Where $\boldsymbol{\mu}$ is the angular moment, \mathbf{r} is the radius, ρ is density, and \mathbf{u} is velocity.

Again, substituting in the analogous variables, one obtains a formula for this torque on the local moment:

$$\boldsymbol{\tau} = \left| \boldsymbol{\mu} \times \bar{\boldsymbol{\omega}} \right| \quad (4)$$

While there were reservations and questions regarding the use of a variable point of reference for the local angular momentum calculation, there was overwhelming agreement with the Lamb vector formulation (Fig. 3). Given this information, plus given the calculated curvature radius near the tornado strongly agreed with the

‘actual’ radius (~ 500m) we have confidence that the data are meaningful.

We put this type of interaction in the context of vortex loops generated by a pulsing rear-flanking downdraft near a mesocyclone. The gyroscopic torque preferentially rotates the vortex loop into the ‘correct’ configuration, where this vorticity reconnects with the mesocyclone and brings the funnel to the ground (Fig. 3, right side).

In addition, we also performed idealized vortex interaction simulations. We performed a balanced initialization of a horizontal vortex loop embedded within a large vertically oriented Rankine vortex. We also varied the orientation of both the Rankine vortex and the horizontal loop. In our initial findings, we found that the small-scale vortex filaments always tended toward greater alignment with the background vorticity field. An example of this interaction is given in Figure 4. Here, an idealized loop (with the curl of the vorticity pointed downward), similar to that generated by a rear flanking downdraft loop, is intersecting a larger, tornado-like Rankine vortex. The intersected portion of the loop immediately aligns, even without the presence of an updraft.

The units of this torque are m^2s^{-2} . Since the torque is acting on the entire cross-section of the angular moment, the total “aligning” on the vortex should be scaled by the area. When examining the units after this scaling, one finds this would be the equivalent of vorticity tendency (s^{-2}).

3.2 Merging parameter

Once the surrounding vortices are aligned with the parent vortex, another mechanism is needed to draw the small-scale vortex upgradient. This “vortex beta” parameter is shown in Figure 5, where an isosurface of 3-D vorticity magnitude (the main feature is the tornado) is colored by the intensity of this vortex-beta parameter. Yellow values indicate high levels of the vortex-beta force.

This vortex-beta force competes against the so-called Magnus effect, which acts through the Lamb-vector term using the mean velocity and the perturbation vorticity:

$$\frac{\partial \mathbf{u}'}{\partial t} \approx -(\boldsymbol{\omega}' \times \bar{\mathbf{u}}) \equiv -\mathbf{I}'_b \quad (5)$$

This force will act to *eject* like-signed embedded vortices *away* from the center of large-scale rotation. Thus, there will be a *competition* between the vortex beta force (strongly dependent upon the *gradient* of like-signed large-scale vorticity) and the Magnus effect in the final merger process. These two forces can define a fundamental balance relationship, and formulated into a “net merging force” :

$$\boldsymbol{\alpha} = \beta + \mathbf{I}'_b = \nabla(\boldsymbol{\mu} \cdot \bar{\boldsymbol{\omega}}) + (\boldsymbol{\omega}' \times \bar{\mathbf{u}}) \quad (6)$$

This upgradient force on like-signed vortical structure is referenced in several places in the literature,

(e.g. *Schechter and Dubin, 1999*), where it is argued that it is energetically favorable for 'prograde' (like-signed) vorticity to be drawn upgradient, while expelling 'adverse' vortical structures.

The units of this force are ms^{-2} . Since the force is acting on the entire cross-section of the angular moment, the total "merging force" on the vortex should be scaled by the area. When examining the units after this scaling, one finds this would be the equivalent of vorticity gradient tendency ($\text{m}^{-1}\text{s}^{-2}$).

4. CONCLUSION AND FUTURE WORK

We have illustrated the concept of utilizing the EM-HD analogy in terms of vortex interaction during tornadogenesis, highlighting two self-organizational mechanisms of reorientation and upgradient advection of local angular moments embedded within a large-scale rotation. Self-alignment and merging of small-scale moments in the presence of a large-scale field is ubiquitous in both EM and HD. It is likely that the reason is linked to the tendency of a physical system to seek the lowest energy state. We plan to investigate other vortex interaction problems (e.g. hurricanes and boundary-layer turbulence) using this "analogous thinking" in upcoming work.

5. REFERENCES

Belevich, M., 2008: Non-relativistic abstract continuum mechanics and its possible physical interpretations. *J.*

Phys. A: Math. Theor. 41 (045401) doi: 10.1088/1751-8113/41/4/045401

Marmanis, H., 1998: Analogy between the Navier-Stokes equations and Maxwell's equations: Application to turbulence. *Phys. Fluids*, 10 (6), 1428-1437.

Martins, A. and Pinheiro, M. J., 2009: Fluidic electrodynamics: Approach to electromagnetic propulsion. *Phys. Fluids* 21, 097103; doi:10.1063/1.3236802

Rotunno, R., 1984: An investigation of a three-dimensional asymmetric vortex. *J. Atm. Sci.*, 41, 283-298.

Schechter, D and Dubin, D, 1999: Vortex motion driven by a background vorticity gradient. *Phys. Rev. Let.* 83 (11) 2919-2194.

Steinhoff and Underhill, 1994: Modification of the Euler equations for "vorticity confinement": Application to the computation of interacting vortex rings, *Phys. Fluids* 6, 2738; doi:10.1063/1.868164 [13]

Thomson, J. J., 1931: On the analogy between the electromagnetic field and a fluid containing a large number of vortex filaments. *Phil. Mag. S. 7.12* (80) 1057-1063

Tripoli, G. J., 1992: A nonhydrostatic numerical model designed to simulate scale interaction, *Mon. Weather Rev.*, 120, 1342-135

Turbulent hydrodynamics	Electromagnetism	Analogous Variables	
Navier-Stokes $\frac{\partial \mathbf{u}}{\partial t} = -(\boldsymbol{\omega} \times \mathbf{u}) - \nabla \left(\frac{p}{\rho} + \frac{u^2}{2} \right) + \nu \nabla^2 \mathbf{u}$	$\frac{\partial \mathbf{A}}{\partial t} = -\mathbf{E} - \nabla \phi$ $\mathbf{E} = \text{electric field}$	Vector and scalar potential	\mathbf{u} \mathbf{A} \mathbf{I} \mathbf{E}
Lamb vector and vorticity $\mathbf{l} \equiv (\boldsymbol{\omega} \times \mathbf{u}) \quad \nabla \cdot \boldsymbol{\omega} = 0 \quad \boldsymbol{\omega} = \nabla \times \mathbf{u}$	$\mathbf{B} = \nabla \times \mathbf{A} \quad \nabla \cdot \mathbf{B} = 0$ magnetic induction Coloumb Thomson		$\boldsymbol{\omega}$ \mathbf{B}
vorticity tendency $\frac{\partial \boldsymbol{\omega}}{\partial t} = -\nabla \times \mathbf{l}' + \nu \nabla^2 \boldsymbol{\omega}$ <small>- viscous term removed</small>	$\frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E}$ Faraday's Law		
Lamb vector tendency and turbulent current (\mathbf{j}) $\frac{\partial \mathbf{l}}{\partial t} = \nabla \times \boldsymbol{\eta} - \mathbf{j}$	$\frac{\partial (\epsilon_0 \mathbf{E})}{\partial t} = c^2 \nabla \times \mathbf{H} - \mu_0 \mathbf{J}$ (zero polarization) Ampere's Law		$\boldsymbol{\eta}$ \mathbf{H}
vorticity field strength ($\boldsymbol{\eta}$) and magnetization $\boldsymbol{\eta} = \mathbf{u}^2 \boldsymbol{\omega} - \mathbf{M}$ $\mathbf{M} = \mathbf{u}(\mathbf{u} \cdot \boldsymbol{\omega}) + \nu \nabla^2 \mathbf{u}$	$\mathbf{H} = \frac{1}{\mu_0} \mathbf{B} - \mathbf{M}$ magnetic field strength and magnetization		
turbulent charge density $\nabla \cdot \mathbf{l} = \mathbf{u} \cdot \nabla \times \boldsymbol{\omega} - \boldsymbol{\omega}^2 \equiv \rho_n$	$\nabla \cdot (\epsilon_0 \mathbf{E}) = \rho_e$ electric charge density		ρ_n ρ_e

Figure 1. A table showing the analogous mathematical structure and variables between the equations of fluid dynamics and electromagnetism. Variables are color-coded to highlight the analogies.




Figure 2. The left side of the diagram shows the geometric configuration of a segment of perturbation vorticity ($\boldsymbol{\omega}'$) embedded within a larger scale vortical field ($\bar{\boldsymbol{\omega}}$). The curl of the second term in the Lamb vector decomposition ($(\bar{\boldsymbol{\omega}} \times \mathbf{u}') \equiv \mathbf{I}'_a$) describes the same type of “counter-torque” yielding alignment of a flywheel in a gimbal gyroscope. This quantity lies in the direction of the cross product of the local moment and the mean rotation, mathematically identical in form to the torque on a current loop or magnetic dipole placed in an external magnetic field (right side of diagram).




Figure 3. There is remarkable comparison of the EM-formulated gyroscopic torque (left) and the Lamb vector formulation (right). The bright yellow patches indicate strong torque. The area in the southeast part of the domain is the region of strong vorticity associated with the advancing gust front associated with the rear flanking downdraft. Also note the reduction of numerical noise in the EM-formulation. In the far right portion of the figure, a schematic of the tornadic vortex evolution is given. For example, a new downdraft pulse is also forming north of the mesocyclone.




Figure 4. Idealized vortex interaction between horizontal vorticity loop (blue) and vertical vorticity Rankine vortex (red). The curl of the vorticity in the loop is downward (opposite the vorticity in the Rankine vortex). Note how the embedded portion of the horizontal loop aligns with the Rankine vorticity.




Figure 5. Isosurface of vorticity magnitude, colored by the magnitude of the “3-D vortex beta vector”. Orange arrow indicates direction of vortex beta force.