

# Sea Surface Pressure Fields from NASA's SeaWinds Scatterometer and Their Impact in NWP

Shannon R. Davis, Mark A. Bourassa, R. Atlas, J. Ardizzone, Eugenia Brin  
James J. O'Brien, David F. Zierden

## 1. Introduction

Spaceborne scatterometers provide marine surface wind vector measurements with unprecedented coverage in space and time. As a result, these instruments can potentially serve as invaluable tools in operational meteorology as well as in the realm of numerical weather prediction. The most recently launched satellite scatterometer,

SeaWinds, relays data from over 94% of the planet's ice-free oceans in a 24-hour period. Here the impact of assimilated SeaWinds data upon a numerical weather models performance is examined in some detail with particular emphasis devoted to the impact of assimilated scatterometer-derived sea surface pressures. Several experiments are conducted using a new NASA/NCAR finite volume model to separately assess the impact of assimilated sea-surface pressure fields derived from SeaWinds and assimilated SeaWinds winds themselves. The results from these experiments are objectively analyzed and the individual impacts of assimilated SeaWinds variables is compared.

## 2. Background

This study involving the assessment impact of assimilated scatterometer-derived pressures in addition to assimilated scatterometer winds is a significant departure from many other earlier works. With preliminary impact studies made by Baker et al. (1984) and Duffy et al (1984), it was established that satellite scatterometer wind data (SeaSAT) alone could provide significant improvements in the surface analyses for major synoptic events. However, less of an impact on the analyses and forecasts for the upper atmospheric levels was realized. In subsequent studies, several initiatives including

those by Duffy and Atlas (1986) and more recently by Atlas(1999,2001), demonstrated that the impact of scatterometer surface winds from could be extrapolated vertically by adjusting the mass of upper levels relative to the adjusted features observed at the surface. Here, an alternative technique to enhancement of scatterometer data impact is advanced. Sea-surface pressures derived from the scatterometer data are assimilated into the model in place of the wind data itself. Assuming a relatively hydrostatic state in the atmosphere, surface pressures represent a three dimensional column of the atmosphere rather than the two-dimensional field depicted by the wind vectors alone. Surface pressure fields additionally affect the mass fields of the atmosphere directly and thus the impact of their assimilation into the model should be much stronger than that of the winds alone. With the use of pressures, many issues regarding complicated boundary layer physics can be circumvented and so too the need for any vertical extrapolation scheme to transmit the influence of the scatterometer data to upper atmospheric levels.

The calculation of sea-surface pressures from the scatterometer wind vectors in this study is made following the works of Harlan O'Brien (1986), Brown and Zeng (1994) and Zierden et al. (2001). Wind vectors from the SeaWinds instrument are produced along a regularly gridded swath that flows. From these wind vectors observations, relative vorticity is determined following a centered difference scheme and blended via a variational technique with a geostrophic vorticity derived from an initial guess/model pressure field. A new surface pressure field is solved numerically using a method of successive overrelaxation.

Figure 1. Coverage for one day from NASA's SeaWinds scatterometer





Figure 2. Experimental Methodology



### 3. Data & Instrument

The SeaWinds instrument was launched by NASA in 1999 aboard the sun-synchronous, polar-orbiting satellite, QuikSCAT. It is a Ku-band scatterometer in the tradition of its predecessors SeaSAT and NSCAT, and functions as an active microwave sensor operating at 13.2 GHz. Unlike its predecessors however, SeaWinds employs twin conically rotating pencil beams in its design, thereby improving the quantity and quality of its observations. With an orbital period of 101 minutes, the coverage of SeaWinds is typically 94% of the ice free oceans in one day. This can be seen graphically in Figure 1.

Since its launch several different data sets have been made available to researchers and operational weather centers by NASA-JPL and NOAA-NESDIS. These include processed science quality and near-realtime wind vectors. Both are employed in this study to be consistent with previous impact studies and to suggest the operational viability of using the SeaWinds winds and wind-derived surface pressures.

### 4. Model

A new Physical-space/Finite Volume global circulation model was employed in the experiments here. This model was developed jointly by NCAR and the NASA Data Assimilation Office. Detailed information about the model is available on the web at: <http://dao.gsfc.nasa.gov/pages/atbd.html>

### 5. Methodology

A series of individual experiments are conducted to assess and compare the impact of assimilated SeaWinds-derived pressures and assimilated SeaWinds surface wind vectors. The methodology for this impact study follows most closely that established by Atlas et al (2001) and is outlined graphically in Figure 2.

### 6. Results & Conclusions

Results and conclusions will be presented at the conference and made available via personal correspondence with the corresponding author

### 7. Acknowledgements

SeaWinds Data used in this experiment was processed and made available by several essential research groups. NASA support was directed from the Ocean Vector Wind Science Team. Data sets were produced by Remote Sensing Systems and sponsored in part by NASA's Earth Science Information Partnership, an association of technical information sites for the Earth sciences and by the NOAA/NASA pathfinder Program for early EOS products; Principal investigator: Frank Wentz. The Center for Ocean Atmospheric Prediction Studies receive its base funding from ONR's Secretary of the Navy grant.

### 8. References

- Atlas, R., 1997: Atmospheric observations and experiments to assess their usefulness in data assimilation. *J. Meteor. Soc. Japan*, **75(1B)**,111-130.
- Atlas, R., S. C. Bloom, R. N. Hoffman, E. Brin, J. Ardizzone, J. Terry, D. Bungato, and J. C. Jusem, 1999: Geophysical validation of NSCAT winds using atmospheric data and analyses. *J. Geophys. Res.*, **104(C5)**,11405-11424.
- Atlas, R. and R.N. Hoffman. 2001: The use of satellite surface wind data to improve weather analysis and forecasting. in *Satellites and Society*, Elsevier, 57-79.
- Baker, W.E., R. Atlas, E. Kalnay, M. Halem, P. M. Woiceshyn, S. Peteherych, and D. Edelmann, 1984: Large-scale analysis and forecast experiments with wind data from the Seasat-A scatterometer. *J. Geophys. Res.*, **89(D3)**,4927-4936.

- T. Liu, and J. J. O'Brien, 1997: Wind observations from new satellite and research vessels agree. *Trans. Amer. Geophys. Union*, **78(51)**, 597-602.
- Bourassa, M.A., D. M. Legler, J.J. O'Brien, and S. Smith, 2001: SeaWinds validation with research vessels. *J. Geophys. Res.*, submitted.
- Brown, R.A., and G. Levy, 1986: Ocean surface pressure fields from satellite sensed winds. *Mon. Wea. Rev.*, **114**, 2197-2206.
- Brown, R.A., and L. Zeng, 1994: Estimating central pressures of oceanic mid-latitude cyclones. *J. Appl. Meteor.*, **33**, 1088-1095.
- Cane, M.A., V. J. Cardone, M. Halem, and I. Halberstam, 1981: On the sensitivity of numerical weather prediction to remotely sensed marine surface wind data: A simulation study. *J. Geophys. Res.*, **86(C9)**, 8093-8106.
- Harlan, J. Jr., and J.J. O'Brien, 1986: Assimilation of scatterometer winds into surface pressure fields using a variational method. *J. Geophys. Res.*, **91**, 7816-7836.
- Bourassa, M.A., M. H. Freilich, D. M. Legler, W. Hoffman, R.N., 1993: A preliminary study of the impact of ERS-1 C-band scatterometer wind data on the European Center for Medium Range Forecasts global data assimilation system. *J. Geophys. Res.*, **98**, 10233-10244.
- Liu, W.T., W.Q. Tang, and P.S. Polito, 1998: NASA scatterometer provides global ocean-surface wind fields with more structures than numerical weather prediction. *Geophys. Res. Lett.*, **25**, 761-764.
- Zierden, D. F., Bourassa, M.A., and J.J. O'Brien, 2000: Cyclone surface pressure fields and frontogenesis from NASA scatterometer (NSCAT) winds. *J. Geophys. Res.*, **105**, 23967-23981.

