

16C.2 LONG-TERM CARBON AND ENERGY FLUXES FOR AN OLD-GROWTH RAINFOREST

Matthias Falk*, Kyaw Tha Paw U and Matt Schroeder
University of California, Davis, Davis, CA 95616

1. INTRODUCTION

Eddy-covariance and biometeorological methods show significant net uptake by an old-growth Douglas-fir forest in southern Washington state, the oldest forest ecosystem (500 years old) in the AmeriFlux network. Annual net ecosystem exchange ($1.3 - 2 \text{ tC ha}^{-1} \text{ yr}^{-1}$) was comparable to younger ecosystems at the same latitude, as quantified in the AmeriFlux program. Data acquisition at the Wind River Canopy Crane research Facility (WRCCRF) site is ongoing since 1997/98. A detailed analysis of the ecosystems response to changing climatic conditions will be presented. Quantities of interest are the carbon flux and water vapor flux between the forest ecosystem and the atmosphere in dependence of environmental variables such as soil and canopy temperature, soil moisture, vapor pressure deficit and photosynthetic active radiation (PAR).




Figure 1: CO_2 -Flux as fitted function of Photosynthetically Active Radiation (PAR) sorted by temperature class for April 1999. Optimal temperature range for ecosystem photosynthesis lies between 5-10 °C.

2. RESULTS AND DISCUSSION

The data set into several subcategories for inter-annual comparison: (1) Summer Drought (June - August) and loss of CO_2 to atmosphere; (2) Main Uptake of CO_2 (March - May) during Spring; (3) Steady State of CO_2 budget in the Winter (December - February) and (4) Fall recharge period (September - November).

* Corresponding author address: Matthias Falk, Univ. of California, L.A.W.R., Davis, CA 95616; email: mbfalk@ucdavis.edu




Figure 2: CO_2 -Flux as a function of PAR and Vapor Pressure Deficit (VPD) for all 1998 data, when VPD reached large values due to 78 day drought. CO_2 -Flux drops off rapidly with VPD.

The actual temporal boundaries between the different categories are not static but depend on climatic factors. The main carbon dioxide uptake period for example can extend well into June and the beginning of July if rains continue to fall and temperatures stay cool. Summers can either be hot and dry (1998) or wet and relatively cool (1999).




Figure 3: Cumulative CO_2 -Flux for Winter and Spring for 1999-2001. Late Spring '99 was cooler and net ecosystem uptake continued through July. In 2000 the uptake slowed significantly in May resulting in a shortfall of 0.7 tC ha^{-1} . Nighttime data was replaced by temperature modeled respiration when $u^* < 0.5 \text{ m/s}$.




Figure 4: Cumulative CO_2 -Flux for Winter and Spring for 1998-2001. The El Nino Summer '98 was extremely dry and warm compared to the following year. Respiration was significantly enhanced. Ecosystem Net Loss was higher by over 0.5 tC ha^{-1} .




Figure 5: Cumulative CO_2 -Flux for 3 1/2 years, 1998-2001. Net Carbon Uptake occurs during times of maximum precipitation and low temperatures from late winter to early summer. Annual NEE is varies from a strong uptake of $2.0 \text{ tC ha}^{-1} \text{ year}^{-1}$ to a net loss of $0.5 \text{ tC ha}^{-1} \text{ year}^{-1}$.

The year 1998 was an El Nino period, where conditions were warmer and drier than normal. By September 1998, soils were at their driest on record, and air temperature and atmospheric vapor pressure deficit

(VPD) were both large. In contrast, temperatures and precipitation in 1999 were more typical of long-term means and soil moisture was larger than for corresponding summer months in 1998. It is important to note that this coniferous ecosystem has a very different seasonality than eastern hardwood and coniferous forests.

	NEE	GEP	TER	TSR	TSR/TER
1998/99	1.5	15.5	14.0	10.9	78 %
1999	2.1	15.2	13.0	8.7	67 %
2000	0.7	16.8	16.3	11.2	69 %
2001	-0.5	--	--	--	--

Table 1: Wind River Ecosystem Exchange Components (All units are in $\text{tC ha}^{-1} \text{ yr}^{-1}$). NEE: Net Ecosystem Exchange, GEP: Gross Ecosystem Productivity, TER: Total Ecosystem Respiration, TSR: Total Soil Respiration. NEE as measured by Eddy covariance at 70m, TSR as measured by Eddy Covariance at 3m and TSR calculated from a respiration temperature model derived from nighttime EC 70m data for $u^* > 0.5 \text{ m/s}$.

3. CONCLUSION

Extent and severity of the summer drought as well as spring temperature and precipitation regime are the main factors in determining respiration, evapotranspiration and photosynthetic uptake and hence ecosystem exchange with the atmosphere.

4. ACKNOWLEDGEMENTS:

This research was funded by the Western Regional Center (WESTGEC) of the National Institute for Global Environmental Change (NIGEC) through the U.S. Department of Energy (Cooperative Agreement No. DE-FC03-90ER61010).

The WRCCRF is operated under joint sponsorship of the University of Washington and the USDA Forest Service/PNW Station and we acknowledge both for significant support. Any opinions, findings and conclusions or recommendations expressed herein are those of the authors and do not necessarily reflect the view of the DOE.