

Jee-Hoon Jeong* , Chang-Hoi Ho, Dao-Yi Gong
Seoul National University, Seoul, Korea

1. Introduction

In recent years, there has been great interest in the Arctic Oscillation (AO), which was introduced as an annular mode of atmospheric circulation by Thompson and Wallace (1998). This primary mode of internal dynamics in the atmosphere predominates the extratropical Northern Hemisphere circulation from the surface to the lower stratosphere and shows an equivalent barotropic structure during cold seasons (November-April) [Thompson and Wallace 2000]. Fluctuations in the AO create a seesaw pattern in which atmospheric pressure and mass in northern polar and mid-latitudes alternate between positive and negative phase [Wallace 2000]. Many studies have indicated that the AO significantly influences on mid- to high latitude climate in Northern Hemisphere [Thompson et al. 2000, Kerr 1999]. For example, the AO-related climate variations involve surface air temperature [Thompson et al. 2000, Rigor et al. 2000], precipitation [Thompson et al. 2000, Cavazos 2000], moisture budget [Boer et al. 2001], sea-ice over north polar and sub-polar regions [Wang and Ikeda 2000], lower tropospheric circulation including the East Asian winter monsoon, Aleutian Low, and Siberian High [Gong et al. 2001, Overland et al. 1999], and extreme climates such as cold waves and blocking activities [Thompson and Wallace 2001].

Although the AO can explain a large portion of the total variance in the atmospheric circulation during warm seasons (May-October) [Thompson and Wallace 2000], most previous studies, however, have emphasized the simultaneous climate influence in cold seasons when the AO is most active. Here we present evidence showing that there is a significant connection between late spring AO and the East Asian summer precipitation on the interannual time scale for the last one hundred year (1899-1999). We use the long-term precipitation record observed in China and surrounding regions.

2. Data preparation

AO indices are determined by time series of the leading principal component of monthly mean sea level pressure in Northern Hemisphere (poleward of 20°N) [Thompson and Wallace 1998]. There are two kinds of AO index. One has a longer time series for

the period 1899-1997, which is derived from observed sea level pressure. The other is available for the period 1958-1999, which is calculated using reanalyzed sea level pressure of the National Centers for Environmental Prediction/the National Center for Atmospheric Research (NCEP/NCAR). Here we use the combined series for period 1899-1999 (prior to 1957 based on the observed data and thereafter on the NCEP/NCAR reanalysis data). The data are prepared and provided by David W. J. Thompson and available via internet at <http://horizon.atmos.colostate.edu/ao>.

Rainfall data at ten stations are selected to investigate the relationship between the AO index and the East Asian summer monsoon precipitation. These stations are located in central China and southern Japan. Time series of seasonal rainfall at the six Chinese stations are available since 1880 [Wang et al. 2000]. More information about these data can also be found in Qian and Zhu (2001). Monthly rainfall data at the four Japanese stations are taken from the Global Historical Climatology Network [Vose et al. 1992]. The rainfall data at the ten stations represent the *Meiyu* rainfall features faithfully due to the high consistent variations of summer rainfall over the region as revealed in the empirical orthogonal function analysis [see Nitta and Hu 1996]. A historical monthly precipitation dataset for global land areas from 1900 to 1998 is also used here, which is gridded at a 5° latitude by 5° longitude resolution [Hulme 1996].

The 200 hPa zonal wind data as well as wind velocities at other levels are taken from the NCEP/NCAR data set for the period 1958-1999 [Kalnay et al. 1996]. The NCEP/NCAR geopotential heights in the troposphere are also utilized.

Since there is strong interdecadal variability in the summer monsoon rainfall and circulation due to the tropical sea surface temperature anomalies [e.g., Hu 1997, Weng et al. 1999], a high-pass filter is necessary to remove the possible influence of these low-frequency variations and get a robust signal at interannual time scale. Thus, a 9-point Gaussian digital filter with weights of 0.01, 0.05, 0.12, 0.20, 0.24, 0.20, 0.12, 0.05 and 0.01 is applied to all data sets. This filter removes variations longer than 10 years and keeps interannual variations only.

3. Summer rainfall

Summer (June, July and August) rainfall is one of the most important indicators of the East Asian summer monsoon. It is widely recognized that the main rainfall band, known as *Meiyu*, is most prominent from the Yangtze River valley to the

* Corresponding author address: Jee-Hoon Jeong, School of Earth and Environmental Sciences, Seoul National University, Seoul, 151-742, Korea, Tel. +82-2-880-8861, Fax. +82-2-876-6795, e-mail: jeehoon@cpl.snu.ac.kr

southern Japan resulting in the east-west directed pattern. Consequently, the long-term mean of summer precipitation shows that 400-500 mm isohyets extend along the Yangtze River, across southern Japan, to the western North Pacific at about 160°E. Over this region, summer rainfall accounts for 40-50% of the annual total precipitation amount [Lau *et al.* 1988, Tao and Chen 1987, Ding 1992]. Thus, the strong year-to-year variation of the summer monsoon is of great interest since it is usually responsible for the severe floods and droughts over the regions.

Our analysis found that the interannual variability in summer rainfall shows an evident connection to the late spring AO index. In particular, the May AO shows the most significant impact on the subsequent summer rainfall. Figure 1 exhibits the geographical distribution of rainfall changes associated with the May AO, shown as the regression coefficients upon the AO index. Here the AO index is normalized before regressing. As shown in the figure, the AO-related summer monsoon rainfall changes show a well-defined large-scale pattern with a strong east-west direction. With the one standard deviation higher index, there are about 20-40 mm decrease in precipitation over the region extending from the Yangtze River valley to the southern Japan and 10-30 mm increase over the southern China.




Figure 1. Changes in summer precipitation (mm) corresponding to a one standard deviation of the May AO index for the period 1900-1998. Grey squares indicate the grids with data availability above 95% in the entire period. Regions above 95% confidence level are shaded. Ten stations are shown as filled circles. Contour interval is 10 mm. Zero contours are omitted.

Rainfall at each station as well as their mean is also analyzed for comparison with the gridded data sets. Correlation coefficients vary from about -0.20 to -0.42. The corresponding precipitation changes are on average -48.3 mm per one standard deviation of the AO index, which are a little larger than the gridded data due to the higher variance of the individual station records. A comparison of correlation coefficients of unfiltered and filtered data also confirms that the strong relationship is attributable mostly to

the interannual covariance.

As the individual station's rainfall is usually influenced by many local environmental factors and thus the relation signal between AO and precipitation could be distorted to some degree. This shortcoming will be improved if the regional mean series are used. In fact, the ten stations' mean rainfall shows a much stronger connection to the AO (see Fig. 2). Some anomalous rainfall events, such as the strong positive anomalies in years of 1954, 1969, 1980, and 1993 are outstanding. Impressively, these events are well correlated with strong negative-phase of May AO. This kind of out-of-phase relationship between AO and rainfall is most predominant in late five decades. For the whole period (93 years), the two timeseries are correlated at the coefficient of -0.45. Given the fact that all data are high-pass filtered, the degrees of freedom should be not much less than 90. Autocorrelation of AO also confirms that. Thus, the above correlation is highly significant at the 99% confidence level.




Figure 2. Time-series of May AO index and summer monsoon rainfall over Yangtze River valley and southern Japan. Shown as the results from a high-pass filter, through which the variations longer than 10 years are removed.

We selected these ten stations based on Fig. 1, as anomaly center is located over the Yangtze River to southern Japan, over there only six stations with data available for period 1880-1999 in eastern China. In southern Japan there are some long-term observations. In fact, some other stations also show significant correlation coefficient (r) for the same period, for example, Nagasaki (129.9°E, 32.7°N), $r = -0.30$ and -0.21 , Miyazaki (131.4°E, 31.9°N), $r = -0.25$ and -0.15 , and Shionomisaki (135.8°E, 33.5°N, data available for 1913-1999), $r = -0.31$ and -0.20 , for filtered and unfiltered series, respectively. In this study we pick out four evenly distributed stations in order to appropriately cover the *Meiyu* region. In addition, 33 stations [Wang *et al.* 2000] scattered over the eastern China with long-term record (1800-1999) are also examined. The results of regression analysis display the similar pattern as shown in Fig. 1 (figure not shown).

4. East Asian summer monsoon circulation

There are at least two major large-scale

circulation systems that dominate the East Asia summer monsoon. One is the East Asian westerly jet stream and the other is the western Pacific subtropical high. Previous studies indicated that the north-south displacement of the jet plays an important role in determining the monsoon onset and retreat. Furthermore, the location of the jet stream is closely related with the fronts that produce an extensive summer monsoon rainfall (*Meiyu*), and thus can be used as a representative circulation for the summer monsoon.

To investigate the associated features in atmospheric circulation with AO, the high-pass filtered summer mean zonal winds at 200hPa are regressed upon the late (May) AO index and spring (March-May) AO index, respectively. Covariance of summer zonal wind with the late spring AO index shows a north-south movement of the middle latitude zonal jet at many latitudes in the Northern Hemisphere (Fig. 3a). The covariance over East Asia is more remarkable than other regions. The evident west-east oriented bands over East Asia are of particular interest since this statistically significant pattern is the strongest anomaly center in the entire Northern Hemisphere. With a positive phase of the AO, stronger westerly wind appears in regions of 40°-50°N and 15°-25°N. Latitudes between these two westerly zones are dominated by strong easterlies. This feature indicates that the fluctuation of AO in late spring is associated with a strong meridionally oriented trough-ridge system spanning the coastal region of East Asia and western North Pacific in summer. This also implies that the anomalous East Asian jet associated with the AO undergoes meridional displacement. The climatological location of the East Asian jet core is around 40°N in summer. For the high positive phase AO years, the jet is displaced poleward and an anomalous westerly center appears at about 45°N, five degrees northward of its normal position. Compared to the late spring AO, the spring AO index shows little weaker correlation with the zonal wind in mid-latitudes over East Asia (figure not shown).

Figure 3b shows the zonal wind changes that are associated with the summer rainfall fluctuation (ten stations' mean series) on the interannual time scale. Obviously, there is a remarkable similarity between the two figures (Figs. 3a versus 3b). Both show similar patterns and indicate a consistent northward shift of the East Asian jet stream. That suggests the importance of the upper tropospheric zonal flow in connecting the late spring AO and the summer rainfall.




Figure 3. Changes in summer zonal wind (m s^{-1}) at 200 hPa associated with a one standard deviation anomaly in the May AO index (a) and in the simultaneous ten-station-mean rainfall for summer (b) for the period 1958-1999. To facilitate comparison rainfall records are multiplied by -1. The contour interval is 0.5 m s^{-1} per standard deviation of May AO and summer rainfall, respectively. Zero contours are omitted. Regions below -0.5 m s^{-1} and above $+0.5 \text{ m s}^{-1}$ are shaded.

In the latitude-height sections, the East Asian summer monsoon is characterized by the multi-cellular meridional circulation over the East Asian sector, extending from the tropics to the mid-latitudes. This multi-cellular structure is referred to as the dominant meridional characteristics of the East Asian summer monsoon. *Lau et al.* (2000) defined the strength of the East Asian summer monsoon as the difference in zonal wind at 200 hPa between average value over the domain of 40°-50°N and 110°-150°E and the domain of 25°-35°N and 110°-150°E. They indicated that the East Asian jet plays an essential role in the dynamical aspect of the summer monsoon. The aforementioned pattern associated with the positive phase of AO is in good agreement with the strong East Asian summer monsoon years (cf. *Lau et al.*'s Fig.4b).

The anomalous zonal mean zonal wind, meridional and vertical wind in the summer over the domain of 110°-150°E are regressed onto the May AO. Results are displayed in Fig. 4. Compared with *Lau et al.*'s Fig.8, the altitude-latitude monsoon structure is evidently similar. Three cells are clearly described. Anomalous easterly with the concurrently strong descending motion are predominant around 30°N. The strong westerly to the north and the relatively weaker one to the south are also significant in the upper troposphere, whereas the contours extend to the lower troposphere. The wind anomaly structure generally shows an equivalent barotropic oscillation with the largest covariance at ~200 hPa. Clearly, the north-south displacement of the zonal wind appears at almost all pressure levels in the

troposphere and is more evident in the upper troposphere above 500 hPa.




Figure 4. Cross section of the zonal mean zonal wind (u), meridional wind (v) and vertical motion (ω) over East Asia (110-150E) regressed onto the May AO index. The u is shown as the contours with interval of 0.3 m s^{-1} . Zero lines are bold and easterly are shown in dashed lines. Regions above 95% confidence level are shaded. The covariance of v and ω are shown as vectors. Values are m s^{-1} (for u and v) and hPa s^{-1} (for ω), corresponding to a one standard deviation anomaly in the May AO index. Before regression the AO index are normalized. The values of largest vectors are 0.47 m s^{-1} for v and $4.3 \times 10^{-3} \text{ hPa s}^{-1}$ for ω .

5. Discussion

The mechanisms that are responsible for the AO-related monsoon variability are still an open question. Some recent results show that the AO signal can propagate downward from stratosphere to troposphere. However, the propagation usually appears in cold seasons when two layers are coupled [Baldwin and Dunkerton 2001]. Anomalies in lower boundary conditions in the mid- and high latitudes also would impact the monsoon system. *Lau et al.* (2000) indicated that the variability of the summer time East Asian jet is only weakly related with the tropical sea surface temperature anomalies. On the other hand, previous results show that there is a significant influence of Asian continental land surface condition on the East Asian summer monsoon [e.g. *Yang and Lau* 1998]. Preliminary result of correlation coefficient between the May AO and regional averaged geopotential height over the domain 110°E - 150°E , 30°N - 40°N in the troposphere shows an upward propagation with time (Fig. 5). This upward propagation of AO signal from the lower troposphere may be related to surface influence. Given the well-known strong influence of AO on the surface climate in mid- and high latitudes in winter and spring, there are possibilities that the AO-related boundary anomalies would in turn impact the atmospheric

circulation and give rise to the anomalous East Asian summer monsoon.

The internal dynamical process in the troposphere may be one very important reason for the relation. The AO is weak in late spring compared to winter, but it is not as weak as indiscernible. It is still the leading mode in inactive seasons. Percentages of variance in monthly mean sea level pressure explained by the AO modes for active and inactive seasons are 21% and 16%, respectively [Thompson and Wallace 2000]. Correlation analysis display that winter AO shows no influence on summer monsoon rainfall. The summer AO shows a considerable correlation to summer rainfall in the concerned region ($r = -0.25$ for period 1899-1999). However, compared to the simultaneous correlation, the AO-leading connection is much stronger (see Table 1). When AO leads by one month, the correlation between May-July AO and summer rainfall is -0.44 . When AO leads by two months, correlation is -0.32 . By three months, correlation down to -0.21 . Obviously, the late spring to early summer AO is most important. The large-scale atmospheric circulation changes in association with the AO also show the consistent patterns in East Asia. That means the cause-and-effect experiences in one to two months. In this time scale, dynamic process, including the persistency (between May and June there is significant correlation in AO indices with a value of 0.30), might be important.

Of course, we need more works on the mechanisms for the details. We are investigating aforementioned possible mechanisms using a general circulation model. Here we primarily address the significant relationship between AO and summer monsoon rainfall and the associated circulation features.

In conclusion, our observations reveal that there is a significant influence of late spring AO on the East Asian summer monsoon circulation and rainfall. This precursory signal will be helpful for us to predict summertime rainfall over East Asia.




Figure 5. Cross correlation between the May AO index and the regional averaged geopotential height over the domain 100°E - 160°E and 30°N - 40°N . Values exceeding the 95% confidence level are shaded.

Table 1. Correlation coefficients between AO index and summer rainfall for the period 1899-1999.

AO	Cor (AO, JJA rainfall)
January	-0.04
February	-0.01
March	-0.10
April	0.06
May	-0.45
June	-0.24
July	-0.08
August	-0.13
Jan.-Mar.	-0.07
Feb.-Apr.	-0.04
Mar.-May	-0.21
Apr.-Jun.	-0.32
May-Jul.	-0.44
Jun.-Aug.	-0.25

References

- Baldwin, M. P., and T. J. Dunkerton, 2001: Stratospheric Harbingers of anomalous weather regimes. *Science*, **294**, 581-584.
- Boer, G. J., S. Fourest, and B. Yu, 2001: The signature of the annular modes in the moisture budget. *J. Climate*, **14**, 3655-3665.
- Cavazos, T., 2000: Using self-organizing maps to investigate extreme climate events: An application to wintertime precipitation in the Balkans. *J. Climate*, **13**, 1718-1732.
- Ding, Y.H. 1992: Summer monsoon rainfalls in China. *J. Meteorol. Soc. Japan*, **70**, 373-396.
- Gong, D. Y., S. W. Wang, and J. H. Zhu, 2001: East Asian winter monsoon and Arctic Oscillation. *Geophys. Res. Lett.*, **28**, 2073-2076.
- Hu, Z. Z., 1997: Interdecadal variability of summer climate over East Asia and its association with 500hPa height and global sea surface temperature. *J. Geophys. Res.*, **102**, 19403-19412.
- Hulme, M., 1992: A 1951-80 global land precipitation climatology for the evaluation of General Circulation Models. *Climate Dynamics*, **7**, 57-72.
- Kalnay, E., M. and co-authors, 1996: The NCEP/NCAR 40-year Reanalysis Project. *Bull. Am. Meteorol. Soc.*, **77**, 437-471.
- Kerr, R. A., 1999: A new force in high-latitude climate. *Science*, **284**, 241-242.
- Lau, K.-M., K.-M. Kim, and S. Yang, 2000: Dynamical and boundary forcing characteristics of regional components of the Asian summer monsoon. *J. Climate*, **13**, 2461-2482.
- Lau, K.-M., G.-J. Yang, and S.-H. Shen, 1988: Seasonal and intraseasonal climatology of summer monsoon rainfall over East Asia. *Mon. Wea. Rev.*, **116**, 18-37.
- Nitta, T., and Z. Z. Hu, 1996: Summer climate variability in China and its association with 500hPa height and tropical convection. *J. Meteor. Soc. Japan*, **74**, 425-445.
- Overland, J. E., J. M. Adams, and N. A. Bond, 1999: Decadal variability of the Aleutian low and its relation to high-latitude circulation. *J. Climate*, **12**, 1542-1548.
- Qian, W. H., and Y. F. Zhu. 2001: Climate change in China from 1880 to 1998 and its impact on the environmental condition. *Climatic Change*, **50**, 419-444.
- Rigor, I. G., R. L. Colony, and S. Martin, 2000: Variations in surface air temperature observations in the Arctic, 1979-97. *J. Climate*, **13**(5), 896-914.
- Tao, S. Y., and L.X. Chen, 1987: A review of recent research of the east Asian summer monsoon in China. In *Monsoon Meteorology*. Eds. C.P. Chang and T. N. Krishnamurti. Oxford University Press, New York., pp60-92.
- Thompson, D. W. J., and J. M. Wallace, 1998: The Arctic Oscillation signature in the wintertime geopotential height and temperature fields. *Geophys. Res. Lett.*, **25**, 1297-1300.
- Thompson, D. W. J., and J. M. Wallace, 2000: Annular modes in the extratropical circulation, Part I: Month-to-month variability. *J. Climate*, **13**, 1000-1016.
- Thompson, D. W. J., and J. M. Wallace, 2001: Regional climate impacts of the Northern Hemisphere annular mode. *Science*, **293**, 85-89.
- Thompson, D. W. J., J. M. Wallace, and G. C. Hegerl, 2000: Annular modes in the extratropical circulation, Part II: Trends. *J. Climate*, **13**, 1018-1036.
- Vose, R. S., R. L. Schmoyer, P. M. Steurer, T. C. Peterson, R. Heim, T. R. Karl, and J. K. Eischeid, 1992: The Global Historical Climatology Network: Long-term monthly temperature, precipitation, sea level pressure, and station pressure data. NDP-041. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Wallace, J. M., 2000: North Atlantic Oscillation/Annular Mode: two paradigms-one phenomenon. *Q. J. Royal Met. Soc.*, **126**, 791-805.
- Wang, S. W. D. Y. Gong, J. L. Ye, and Z. H. Chen, 2000: Seasonal precipitation series over China since 1880. *Acta Geograph. Sinica*, **55**, 281-293. (In Chinese).
- Wang, J., and M. Ikeda, 2000: Arctic Oscillation and Arctic sea-ice oscillation. *Geophys. Res. Lett.*, **27**, 1287-1290.
- Weng, H.-Y., K.-M. Lau, Y. K. Xue, 1999: Multi-scale summer rainfall variability over China and its long-term link to global sea surface temperature variability. *J. Meteor. Soc. Japan*, **77**(4), 845-857.
- Yang, S., and K.-M. Lau, 1998: Influence of sea surface temperature and ground wetness on Asian summer monsoon. *J. Climate*, **11**, 3230-3246.