

## 1.7

### APPLICATION OF INVERSE ANALYSIS FOR SURFACE TEMPERATURE PREDICTION OF SEMI-ARID REGION OF N-E BRAZIL

Romulo da Silveira Paz\*  
Departamento de Ciências Atmosféricas  
Universidade Federal de Campina Grande - UFCG  
Campina Grande-PB, Brazil

Zaqueu Ernesto da Silva  
Departamento de Tecnologia Mecânica  
Universidade Federal da Paraíba - UFPB  
João Pessoa, PB, Brazil

and  
Sukaran Ram Patel  
Departamento de Ciências Atmosféricas  
Universidade Federal de Campina Grande - UFCG  
Campina Grande, PB Brazil

## 1. INTRODUCTION

The earth's surface temperature is an important parameter in the atmospheric phenomena and processes occurring in the boundary layer. It involves in the surface energy balance, evaporation, evapotranspiration and desertification processes and is also considered as an indicator of environmental degradation and climate change. Further, the determination of diurnal variation of ground surface temperature plays a critical role in the modeling of atmospheric planetary boundary layer and, by extension, in mesoscale numerical weather prediction (NWP) models. At the ground surface, the various flux terms of the energy balance equation unfortunately are not, linear function of the surface temperature ( $T_g$ ). These terms include: short wave flux due to solar radiation, infrared flux from the atmosphere, thermal emission at the ground, sensible heat transfer between the air and the ground, latent heat flux due to evaporation/condensation and rainfall, and soil heat flux due to the vertical thermal gradient within the soil. Of these, only the first has well defined, externally forced diurnal and seasonal periods and then only under cloudless skies. Therefore, the difficulty in obtaining an

analytical solution for the heat equation lies not in the differential equation itself but rather in accurately prescribing the external forcing and in the nonlinear feedback nature of the upper boundary condition. For this reason, the value of the surface temperature ( $T_g$ ), in NWP-type models are usually determined by numerical methods. Deardorff (1978) found the so-called force-restore method to be the most satisfactory. This method is used in many NWP-type models (Zhang and Anthes, 1982; Tuccillo and Phillips, 1986). Bhumralkar (1975) has performed numerical experiments on the computation of ground surface temperature in an Atmospheric General Circulation Model. Recently Best (1998) described a model to predict the surface temperature of variety of surfaces, by solving the surface energy balance equation iteratively, using the standard meteorological data. Figueiredo (2000) has used a statistical model to adjust the prognosticated maximum and minimum temperature for 24 hours using the limited area atmospheric model for the city of Botucatu-SP, Brazil. Antonio and França (2000) have estimated the earth's surface temperature using the NOAA-AVHRR satellite data. Peres e Câmara (2002) have studied the estimation of earth's surface temperature using inverse theory and METEOSAT Second Generation (MSG) data, based on the on the method of Watson (1992) and Faysash (1999). In this study the inverse theory is applied for the prediction of the surface temperature of semi-arid region of N-E

---

\*Corresponding author address: Romulo da Silveira Paz, Departamento de Ciências Atmosféricas, Universidade Federal de Campina Grande-PB, 58109-970, Brazil, e-mail [rômulo@camboriu.jpa.com.br](mailto:rômulo@camboriu.jpa.com.br)

Brazil, by solving the surface energy balance equation.

Recently the inverse theory has received much attention in micrometeorological studies and several successful applications have been presented within the last decade (Siquira et al., 2003, 2000; Raupach, 1988; 1989a,b; Denmead and Roupach 1993; Denmead, 2000, Denmead et al., 2000; Katul et al., 1997, 2001; Massman and Weil 1999; Leuning 2000; Leuning et al., Simon, et al., 2002; Hsich, et al., 2003; Paz, 2002; Paz et al., 2004; among others). In short the inverse modeling technique uses a measured data set of an atmospheric quantity and an assumed model relationship that describes the physical processes of the quantity to produce the measured data set as a set of parameters (Wolff and Bange, 2000). In other words the technique uses appropriate model assumptions that are based on theoretical assumptions to fit measured data (Zittel, et al., 2002). The technique is based on the assumption of a relationship (operator **D**) between the model parameter **M** and the measured parameter **C**, i.e.,  $\mathbf{C}_{obs} = \mathbf{D} \mathbf{M}_{mod}$ , where, **D** is also called as dispersion matrix (ixj terms),  $\mathbf{C}_{obs}$  is a vector (with i rows) and  $\mathbf{M}_{obs}$  is a vector with j rows.

## 2. SITE AND INSTRUMENTATION

The experimental site is located in the semi-arid region of the institute-basin of Federal University of Campina Grande-UFCG, in the city of São João do Cariri (latitude 07°22'44"S and longitude 36°32'00"W) and altitude approximately 465 m, covered with sparse vegetation of the type caatinga and pasture. The region is considered to be semi-arid with less abundant rainfall from the month of February to April. The soil is rocky type and the vegetation is sparse and gives the appearance of strips alternatively between the rocky soil and the sparse vegetation. The site location is shown in Figure 1., which contains a map of the Paraíba state of Brazil.

A micrometeorological tower of ten meter height is located in the middle of the sparsely vegetated surface of the experimental site described above. The net radiometer and radiometer (both Campbell Scientific Inc., Q-7) were installed at the canopy top at the height of 6 meter. The soil temperature were measured by the temperature sensor (Campbell Scientific Inc., model 108) of precision of 0,001° C, installed one at the surface (to measure surface temperature) and the other three in the depths of 2, 5 and 15

cms. The air temperatures were measured by, two Cromel Constantin thermocouple of 25 μm and 74 μm diameters, one located at the surface of the vegetation and the other at 10 m height. The temperature and specific humidity were also measured by a HMP 45C (Campbell Scientific Inc.) thermo hygrometer located at the top of the bush. The wind velocity, temperature and specific humidity were measured at 1,5 and 10 m heights. The precipitation and the humidity of the soil were also measured. The observations for this work were made during the period from 28-02-2001 to 09-03-2001 and from 07-11-2001 to 16-11-2001, the rainy and the dry seasons respectively. The measurements were taken during the interval of 20 minutes by the data acquisition system CR23X of Campbell Scientific Inc., continuously connected to the battery of 12 volts and 55 AH, accompanied to a solar panel of potential of 50 W. The datalogger is programmed to control all the experiment and to unload the data to the microcomputer.

## 3. METHODOLOGY AND DISCUSSION

The prognostic equation for the surface temperature,  $T_g$  is obtained in Limited Area Model (ALM) at grid points from the surface energy balance equation suggested by Bhumrarkar (1975) as:

$$G(0)=S - R_L - LE -H \quad (1)$$

Where,

$G(0)$  represents soil heat flux at the surface

$S$  is a solar energy reaching the earth's surface

$R_L$  is net long wave radiation

$LE$  represents latent heat flux and

$H$  represents sensible heat flux

The soil heat flux is defined by

$$G(z, t) = \lambda \frac{\partial T_s(z, t)}{\partial z} \quad (2)$$

where  $\lambda$  is a soil thermal conductivity and  $T_s$  is soil temperature.

$T_s$  may be obtained by solving the soil heat transfer equation for (simplification) homogeneous soil

$$\frac{\partial T_s}{\partial t} = \frac{\lambda}{c_g} \frac{\partial^2 T_s}{\partial z^2} \quad (3)$$

with boundary condition

$$T_s(z, t) = \bar{T} + \Delta T_0 \text{ sen}(\omega t) \quad (4)$$

as,

$$T_s(z, t) = \bar{T} + \Delta T_s e^{-z/D} \text{sen}(\omega t - z/D) \quad (5)$$

where

$c_g$  is soil specific heat ( $\text{J kg}^{-1} \text{K}^{-1}$ )

$\bar{T}$  is daily mean temperature

$\Delta T_0$  is surface thermal amplitude

$\omega$  is frequency of oscillation and

$$D = \left( \frac{2\lambda}{c_g \omega} \right)^{1/2} \text{ is damping depth.}$$

Now substituting (5) in (2) one may have

$$G(z, t) = \left( \frac{\omega c_g \lambda}{2} \right)^{1/2} \left[ \frac{1}{\omega} \frac{\partial T_s(z, t)}{\partial t} + T_s(z, t) - \bar{T} \right] \quad (6)$$

A rate of temporal variation of a layer at depth Z may be written in the form

$$c_g \frac{\partial T_s}{\partial t} = - \left[ \frac{G(Z, t) - G(0, t)}{Z} \right] \quad (7)$$

As, the aim of this work is to calculate the surface temperature of the soil, rather than the mean temperature of soil at finite depth, so eq. (7) may be applied for a layer of soil of 1 cm depth, and considering that the mean temperature of this layer is approximately equal to the surface temperature,  $T_g$ ; so one may have

$$T_s(1, t) \approx T_g(t) \quad (8)$$

Equation (8) represents a reasonable approximation for the present work, seeing the uncertainty of the definition of the surface itself specially for the regional scale. Now from eqs (2), (3) and (4) for the layer of thickness  $Z=0,01$  one may have

$$c_g \frac{\partial T_g}{\partial t} = G(0) - \left( \frac{\omega c_g \lambda}{2} \right) \left[ \frac{1}{\omega} \frac{\partial T_g}{\partial t} + T_g - \bar{T} \right]$$

which can be solved for  $G(0)$  following Bhumhalkar(1975) as:

$$T_g^{t+\Delta t} = T_g + \frac{f_1}{f_2 + f_3} \quad (9)$$

where,

$$f_1 = R_n - LE - H - \left( \frac{\lambda c_g \omega}{2} \right)^{1/2} (T_g - \bar{T}) \quad (10)$$

$$f_2 = \left( \frac{c_1}{\Delta t} \right) + 4\sigma_B T_g^3 \quad (11)$$

$$f_3 = c_H \left( 1 + \frac{L}{c_p} w_g \frac{dq_s}{dT_g} \right) + \left( \frac{\lambda c_g \omega}{2} \right)^{1/2} \quad (12)$$

where other notations are defined in (Bhumhalkar, 1975 and Paz, 2002).

Now the surface temperature can be calculated from eq. (9) where H and LE are obtained from the similarity theory of Monin-Obukhov, using forms of Bussinger et al., (1971) or other formulation for the similarity empirical functions of Monin-Obukhov.

The method of Levenberg-Marquardt (Press et al., 1992) is used to solve the problem of identification. The adjust of the merit function  $S^2$  can be written for the temperature in the form

$$S^2(\beta, t) = \sum_{i=1}^N \left[ \frac{T_{g(\text{obs})}^i - T_{g(\text{mod})}^i}{d_i} \right]^2 \quad (13)$$

The soil's prognostic temperature  $T_g^i$  presents the following functional dependency:

$$T_g^i = F(R_n, T_g, LE, H, t, z, z_0, \lambda, \omega, c_g, K_g, \rho, c, c_p, q, q_s, u) \quad (14)$$

The data obtained were submitted a consistency analysis based on the conventional criteria within the practical norms of the operational meteorology.

The process of the identification requires a preliminary analysis of the sensibility of the variable obtained by model in terms of the parameters and objectives of the estimation. The coefficient of reduced sensibility is used and represented graphically to evaluate the possibility of the satisfactory application of the method of identification. The physical properties of air, soil and wind velocity are evaluated. The analysis of wind velocity although a variable furnished by the micrometeorological station, justifies the analysis because it evaluates indirectly the sensibility of the model to the

Richardson number to include the effect of the stability of the boundary layer. So, finally the one may have

$$T_g^{t+\Delta t} = F(R_n, T_g, t, z_0, \lambda, \bar{\omega}, c_g, c, c_p, u, t) \quad (15)$$

A numerical code is developed in FORTRAN language to perform an analysis for the prognostic model for the surface temperature and to calculate the coefficient of sensibility. In this case the sensibility of the temperature to surface roughness ( $z_0$ ), the frequency of the surface temperature ( $\bar{\omega}$ ), humidity parameter ( $w_g$ ), specific heat of soil ( $c_g$ ), thermal conductivity of the soil ( $\lambda$ ), and specific heat of air ( $c$ ). Further, a sensibility in relation to wind velocity ( $u$ ) considered as parameter in calculating the Richardson number. The surface temperature calculated from the model and the corresponding measured values are shown in figures (1) and (2) for the humid and dry periods, before the identification.




Fig. 1 Observed and modeled values of Surface temperature before the identification for humid period

The performance of the model as seen in the figures are satisfactory, although there are mean relative errors of 12.7% and 8.4% for the humid and dry periods respectively.




Fig. 2 Observed and modeled values of surface temperature before identification for dry period

The figures (3) and (4) represent the modeled and measured values of the surface temperature after the identification of the parameters. The relative mean errors after identification, reduces to 1.4% and 1.1% for humid and dry periods respectively. It can be seen from the figures (3) and (4) that the dispersion of the data has reduced significantly, and there is a quite good agreement between the modeled and measured values after the identification for both, humid and dry periods. This shows the efficiency of the inverse modeling for surface temperature prediction. The further details of this study will be reported in a future paper.




Fig. 3 Observed and modeled values of surface temperature after identification for humid period




Fig. 4 Observed and modeled values of surface temperature after identification for dry period

#### 4. REFERENCES

- Antônio, R. N., and França, J. R. A.; 2002, Estimation of earth's surface temperature using NOAA-AVHRR, Anais do XII Congresso Brasileiro de Meteorologia, pp 2874-2878, Faz do Iguaçu (in portuguese).
- Best, M. J., 1998, A model to predict surface temperatures, *Boundary Layer Meteorol.* 88, 279-306.
- Bhumralkar, C. M., 1975, Numerical experiments on the computation of ground surface temperature in atmospheric general circulation models, *J. Appl. Meteorol.* 14, 1246-1258.
- Deardorff, J. W., (1978) Efficient prediction of ground surface temperature and moisture, with inclusion of a layer of vegetation, *J. Geophys. Res.*, 83, 1889-1903.
- Denmead, O. T., (1995): Novel meteorological methods for measuring trace gas fluxes. *Philos. Trans. R. Soc. London, A* 351, 383-396.
- Denmead,, O. T., Raupach, M. R., (1993): Methods for measuring atmospheric gas transport in agricultural and forest systems, in *Agricultural ecosystem effects on trace gases and global climate change*, Eds. J. M. Duxbury, L. A. Harper, A. R. Mosier and D. E. Rolston, American Society of Agronomy, Madison, USA.
- Denmead, O. T., Harper, . A., and Sharpe, (2000): Identifying sources and sinks os scalars in a corn canopy with inverse Lagrangian dispersion analysis, I. Heat. *Agric. Forest Meteorol.* 108, 67-73.
- Faysash, A., and Smith, E. A., (1999), Simultaneous land surface temperature-emissivity retrieval in the infrared split window, *J. Atmos. Oceanic. Technol*, 16, pp 1673-1689.
- Figueiredo, J. C., (2000), Surface air Temperature prediction of for the city of Botucatu using the data generated by limited area model, Master's dissertation in Agronomy, Unesp/FCT, Botucatu-SP, Brazil, 100p.
- Hsieh, C-I., Siqueira, M., Katul, G., Chu, C.-R., (2003):Predicting scalar source-sink and flux distributions within a Forest canopy using a 2-D Lagtangan stochastic disperson model, *Boundary-Layer Meteorol.* 109, pp. 113-138
- Katul, G. G., Lai; C.-T, Siquira, M., Schafer, K., Albertsson, J., Wesson, K., Ellsworth, D., and Oren, R., (2001) Inferring scalar sources and sinks within canopies using forward and inverse methods. Observations and modeling of land-surface fluxes within hydrological systems, V. L. Albertson and J. D. , eds. American Geophysical union, Washington, D. C., PP. 31-45.
- Katul, G. G., Oren, R., Ellsworth, D., Hsieh, C., Phillips, N., and Lewin, K., (1997): A Lagrangian dispersion model for predicting CO<sub>2</sub> sources and sinks, and fluxes in a uniform loblolly pine (pinusd taeda l.) stand, *J. Geophys. Res.* 102, pp. 9309-9321.
- Leuning, R., (2000) Estimation of scalar source/sink distributions in plant canopies using Lagrangian dispersion analysis: Corrections for atmospheric stability and comparison with a multiplayer canopy model, *Bound.-Layer Meteorol.*, 96, 233-249.

- Leuning, R. O., Denmead, O., Miyatani, A., and Kim, J., (2000): Source/sink distribution of heat water vapor, carbon dioxide and methane in a rice canopy estimated using Lagrangian dispersion analysis, *Agric. Forest Meteorol.*, 104 233-249.
- Paz, R. S., (2002): Inverse analysis applied to Regional Circulation Meteorological Model-MAL Doctorate thesis submitted to Mechanical Engineering Department, Federal University of Paraiba-UFPB (in Portuguese).
- Paz, R. S., Da Silva, Z. E., and Patel, S. R., (2004): Inverse theory applied to surface layer radiation in São João do Cariri-PB N-E Brazil, Eighth symposium on integrated observing and assimilation systems for Atmospheric, Oceans, and land Surface (IOAS-AOLS), American Meteorological Society, Seattle, Washington 11-15 January, 2004, paper number 1.3 (in this issue).
- Peres, L., and Camara, C., (2002), Inversion of the radiative transfer equation for estimation of emissivity and the earth's surface temperature using the satellite data MSG: success and difficulties, *Anais do XII Congresso Brasileiro de Meteorologia*, pp. 3091-3101, Foz de Iguaçu (in Portuguese).
- Press, W. H., Teukolsky, S. A., Vetterling, E., and Flannery, P. B., (1992) *The art of scientific computing*, second edition, Cambridge University Press, New York.
- Raupach, M. R., (1988): Canopy transport processes, pp.95-127, in *Flow and transport in natural environment*, Eds. W. L. Steffen, and O. T. Denmead, Springer-Verlog, New York, USA.
- Raupach, M. R., (1989a): Applying Lagrangian fluid mechanics to infer scalar source distributions in vegetation canopies, *Agric. Forest Meteorol.*, 47, 85-108.
- Raupach, M. R., (1989b): A practical Lagrangian method for relating scalar concentrations to source distributions in vegetation canopies, *Q. J. R. Meteorol. Soc.*, 115, 609-633.
- Simon, E., Ammann, C., Busch, J., Meixner, K., Kesselmeier, J., (2002): Applying Lagrangian dispersion analysis to the exchange of water, sensible heat within a cereal crop canopy: A sensitivity study and comparison with leaf level measurements, 15<sup>th</sup> Symposium on Boundary Layer and Turbulence, 15-19 July 2002, Wageningen, the Netherlands, pp. 535-538
- Siqueira, M., Leuning, R., Kolle, O., Kelliher, F. M., and Katul, G. G., (2003): Modeling sources and sinks of CO<sub>2</sub>, H<sub>2</sub>O and heat within a Siberian pine forest using three inverse methods, *Q. J. R. Meteorol. Soc.*, 129, pp.1373-1393.
- Siqueira, M., Lai, C. T., and Katul, G. G., (2000): Estimating scalar sources, sinks and fluxes in a Forest canopy using Lagrangian, Eulerian, and hybrid inverse model, *J. Geophys. Res.*, 105, 29475-29488.
- Tuccillo, J. J., and Phillips, N. A.; 1986, Modeling of physical processes in Nested Grid Model, Technical procedures bulletin no. 36.3, National Weather Service, Silver Spring, Md.
- Wolff, M. and Bange, J., (2000): Inverse method as an analyzing tool for airborne measurements, *Meteor. Z.*, N. F., 9, PP. 361-376.
- Zhang, D., and Anthes, R. A.; 1982, A high-resolution model of the planetary layer-sensitivity tests and comparison with SESA ME -79 data. *J. Appl. Meteorol.*, 21, 1594-1609
- Zittel, P., Deierling, W. and Babgej J., (2002): Using the inverse method to obtain area averaged turbulent fluxes from airborne measurements at one low altitude, 15<sup>th</sup> Symposium on Boundary Layer and Turbulence, American Meteorological Society, 15-19 July 2002, Wageningen, the Netherlands, pp. 580-581.